

Riordan graphs II: Spectral properties ^{*}

Gi-Sang Cheon[†], Ji-Hwan Jung[†], Sergey Kitaev[‡] and Seyed Ahmad Mojallal[†]

gscheon@skku.edu, jh56k@skku.edu, sergey.kitaev@cis.strath.ac.uk, mojallal@skku.edu

Abstract

The authors of this paper have used the theory of Riordan matrices to introduce the notion of a Riordan graph in [3]. Riordan graphs are proved to have a number of interesting (fractal) properties, and they are a far-reaching generalization of the well known and well studied Pascal graphs and Toeplitz graphs, and also some other families of graphs. The main focus in [3] is the study of structural properties of families of Riordan graphs obtained from certain infinite Riordan graphs.

In this paper, we use a number of results in [3] to study spectral properties of Riordan graphs. Our studies include, but are not limited to the spectral graph invariants for Riordan graphs such as the adjacency eigenvalues, (signless) Laplacian eigenvalues, nullity, positive and negative inertias, and rank. We also study determinants of Riordan graphs, in particular, giving results about determinants of Catalan graphs.

Keywords: Riordan graph, adjacency eigenvalue, Laplacian eigenvalue, signless Laplacian eigenvalue, inertia, nullity, Rayleigh-Ritz quotient, Pascal graph, Catalan graph

2000 Mathematics Subject Classification: 05A15, 05C50

1 Introduction

The authors of this paper have used the theory of *Riordan matrices* to introduce the notion of a *Riordan graph* in [3]. Riordan graphs are a far-reaching generalization of the well known and well studied *Pascal graphs* [9] and *Toeplitz graphs* [12], and also some other families of graphs. The Pascal graphs are constructed using *Pascal's triangle* modulo 2, and Pascal's triangle itself has motivated the appearance of the area of Riordan matrices [17], an active area of research these days. A Toeplitz graph $G = (V, E)$ is a graph with $V = \{1, \dots, n\}$ and $E = \{ij \mid |i - j| \in \{t_1, \dots, t_k\}, 1 \leq t_1 < \dots < t_k \leq n - 1\}$.

Riordan graphs are proved to have a number of interesting (fractal) properties [3], which can be useful, e.g. in creating computer networks [9] with certain desirable features, such as

- the design is to be simple and recursive;
- there must be a universal vertex adjacent to all others;
- there must exist several paths between each pair of vertices.

Also, Riordan graphs can be useful when designing algorithms to compute values of graph invariants.

^{*}This work was supported by the National Research Foundation of Korea(NRF) grant funded by the Korea government(MSIP) (2016R1A5A1008055) and the Ministry of Education (NRF-2016R1A6A3A11930452).

[†]Applied Algebra and Optimization Research Center, Department of Mathematics, Sungkyunkwan University, Suwon 16419, Republic of Korea

[‡]Department of Computer and Information Sciences, University of Strathclyde, 26 Richmond Street, Glasgow, G1 1XH, United Kingdom

The main focus in [3] is the study of *structural properties* of families of Riordan graphs obtained from infinite Riordan graphs, which includes the fundamental *Riordan Graph Decomposition Theorem* (see Theorem 2.4 below) and the generalization of a number of known results for the Pascal graphs.

In this paper, we study *spectral properties* of Riordan graphs. The *spectral graph theory* studies relations between graph properties and the *spectrum* (i.e. the set of eigenvalues) of the adjacency matrix, or (signless) Laplacian matrix, which can be useful in various contexts. For example,

- the *second largest* eigenvalue of a graph gives information about *expansion* and *randomness* properties of a graph;
- the *smallest* eigenvalue gives information about the *independence* and *chromatic numbers* of a graph;
- *interlacing* of eigenvalues (see the definition in Lemma 2.14) gives information about *graph substructures*;
- the fact that eigenvalue multiplicities must be integral provides strong restrictions on graph parameters, e.g. in the case of *strongly-regular graphs*.

In relation to our paper, in [4], several spectral properties of Pascal graphs were studied by exploring their spectral graph invariants such as the algebraic connectivity, the first three largest Laplacian eigenvalues and the nullity. In this paper, we obtain results on several other spectral graph invariants for Pascal graphs. More importantly, we develop the spectral theory for many other classes of Riordan graphs. Our studies include, but are not limited to spectral graph invariants for Riordan graphs such as the *adjacency eigenvalues*, (*signless*) *Laplacian eigenvalues* (see Section 3), positive and negative *inertias* (see Section 6), nullity (see Section 7), and *rank* (see Theorem 7.5). Our key achievement is to give relations for Laplacian spectral radius and algebraic connectivity of Riordan graphs in term of given generating functions. Also, we give a new version of Rayleigh-Ritz quotient for Riordan graphs.

One of the essential difficulties we faced was the fact that the number of edges in Riordan graphs is not known in the general case. This parameter is normally known when dealing with spectral graph invariants. Still, we were able to obtain non-trivial interesting spectral results for various classes of Riordan graphs. In some cases, we use certain structural results in [3] to achieve our goals.

The paper is organized as follows. In Section 2, we give a list of necessary notions, notations and known results. In Section 3, we study the eigenvalues and (signless) Laplacian eigenvalues of Riordan graphs. In Section 4, vertex degrees in Riordan graphs are studied to obtain some bounds for the largest and second smallest Laplacian eigenvalues of Riordan graphs. The *Rayleigh-Ritz quotient* for Riordan graphs is also studied in Section 4. In Section 5, we give results on eigenvalues and Laplacian eigenvalues of Pascal and Catalan graphs. In Section 6, results pertaining to the positive and negative inertias of a Riordan graph, and their complements, are given. In Section 7, we present several results on the nullity and rank of Riordan graphs. Finally, in Section 8, we study determinants of graphs in certain subclasses of Riordan graphs. In particular, we give results on determinants of Catalan graphs.

2 Preliminaries

Graphs in this paper are normally on the vertex set $[n] := \{1, 2, \dots, n\}$. Also, we let $V_o := \{j \in [n] \mid j \text{ is odd}\}$ and $V_e := \{j \in [n] \mid j \text{ is even}\}$. Suppose G is a graph with a vertex set V . For $W \subseteq V$, we denote the subgraph in G induced by W by $\langle W \rangle_G$, or simply by $\langle W \rangle$ if G is understood from the context. The neighborhood of a vertex v , denoted by $N_G(v)$, is the set of vertices in a graph G adjacent to v .

In this paper, the graphs N_n , K_n , $K_{m,n-m}$ ($1 \leq m \leq \lfloor \frac{n}{2} \rfloor$), and P_n are, respectively, the *null graph*, the *complete graph*, the *complete bipartite graph* with parts of sizes m and $n - m$, and the *path graph* on n vertices. Also, $A \equiv B$ means that matrices A and B equal modulo 2. Similarly, $A(i, j) \equiv B(i, j)$ means the (i, j) th element in A equals the (i, j) th element in B modulo 2.

2.1 Riordan graphs

Let $\kappa[[z]]$ be the ring of formal power series over an integral domain κ . If there exists a pair of generating functions $(g, f) \in \kappa[[z]] \times \kappa[[z]]$, $f(0) = 0$ such that for $j \geq 0$,

$$gf^j = \sum_{i \geq 0} \ell_{i,j} z^i,$$

then the matrix $L = [\ell_{i,j}]_{i,j \geq 0}$ is called a *Riordan matrix* (or, a *Riordan array*) over κ generated by g and f . Usually, we write $L = (g, f)$. Since $f(0) = 0$, every Riordan matrix (g, f) is an infinite lower triangular matrix. If a Riordan matrix is invertible, it is called *proper*. Note that (g, f) is invertible if and only if $g(0) \neq 0$, $f(0) = 0$ and $f'(0) \neq 0$.

For a Riordan matrix (g, f) over \mathbb{Z} , the matrix $L = [\ell_{i,j}]_{i,j \geq 0}$ defined by

$$\ell_{i,j} \equiv [z^i]gf^j \pmod{2},$$

is called a *binary Riordan matrix*, and it is denoted by $\mathcal{B}(g, f)$. The *leading principal matrix of order n* in $\mathcal{B}(g, f)$ is denoted by $\mathcal{B}(g, f)_n$.

The following result is well known as the *Fundamental Theorem of Riordan matrices (FTRM)*.

Lemma 2.1 (FTRM, [17]). *Let $R = (g, f)$ be a Riordan matrix and let $RA = B$ where A and B are two infinite vectors with the generating functions $a(z)$ and $b(z)$, respectively. Then $b = ga(f)$.*

The following definition gives the notion of a Riordan graph in both *labelled* and *unlabelled* cases. We note that throughout this paper the graphs are assumed to be *labelled* unless otherwise specified.

Definition 2.2. A simple *labelled* graph G on the vertex set $[n]$ is a *Riordan graph* of order n if the adjacency matrix of G , $\mathcal{A}(G)$, is an $n \times n$ symmetric $(0,1)$ -matrix such that

- its main diagonal entries are all 0, and
- its lower triangular part below the main diagonal is $\mathcal{B}(g, f)_{n-1}$

for some Riordan matrix (g, f) over \mathbb{Z} , i.e., by using Riordan language,

$$\mathcal{A}(G) = \mathcal{B}(zg, f)_n + \mathcal{B}(zg, f)_n^T. \quad (1)$$

We denote such G by $G_n(g, f)$, or simply by G_n when the pair (g, f) is understood from the context, or it is not important. A simple *unlabelled* graph is a *Riordan graph* if at least one of its labelled copies is a Riordan graph.


We note that the choice of the functions g and f in Definition 2.2 may not be unique. If G is a Riordan graph and $\mathcal{A}(G) = [r_{ij}]_{i,j \geq 1}$, then for $i > j \geq 1$,

$$r_{i,j} \equiv [z^{i-1}]zg f^{j-1} \pmod{2} \equiv [z^{i-2}]gf^{j-1} \pmod{2}. \quad (2)$$

For example, the Riordan graph $G_n\left(\frac{1}{1-z}, \frac{z}{1-z}\right)$ is called the *Pascal graph* of order n , and is denoted by PG_n . For another example, the *Catalan graph* $CG_6(C, zC)$ shown below, where

$$C = \frac{1 - \sqrt{1 - 4z}}{2z} = \sum_{n \geq 0} \frac{1}{n+1} \binom{2n}{n} z^n = 1 + z + 2z^2 + 5z^3 + 14z^4 + \dots,$$

is given by the adjacency matrix $\mathcal{A}(CG_6)$:

$$\mathcal{A}(CG_6) = \begin{bmatrix} 0 & 1 & 1 & 0 & 1 & 0 \\ 1 & 0 & 1 & 0 & 1 & 0 \\ 1 & 1 & 0 & 1 & 1 & 1 \\ 0 & 0 & 1 & 0 & 1 & 0 \\ 1 & 1 & 1 & 1 & 0 & 1 \\ 0 & 0 & 1 & 0 & 1 & 0 \end{bmatrix}$$


Definition 2.3. A Riordan graph $G_n(g, f)$ is *proper* if the binary Riordan matrix $\mathcal{B}(g, f)_{n-1}$ is proper. Thus, in a proper Riordan graph a vertex i is adjacent to the vertex $i + 1$ for $i \geq 1$.

If a Riordan graph $G_n(g, f)$ is proper then the Riordan matrix (g, f) is also proper because $g(0) \pmod{2} \equiv f'(0) \pmod{2} \equiv 1$. The converse to this statement is not true. For instance, $(1, 2z + z^2)$ is a proper Riordan matrix but $G_n(1, 2z + z^2)$ is not a proper Riordan graph.

The following theorem about the adjacency matrices of Riordan graphs is a key result in [3].

Theorem 2.4 (Riordan Graph Decomposition, [3]). *Let $G_n = G_n(g, f)$ be a Riordan graph with $[z^1]f = 1$. Then*

- (i) *The adjacency matrix $\mathcal{A}(G_n)$ satisfies*

$$\mathcal{A}(G_n) = P^T \begin{bmatrix} X & B \\ B^T & Y \end{bmatrix} P \quad (3)$$

where $P = [e_1 \mid e_3 \mid \cdots \mid e_{2\lceil n/2 \rceil - 1} \mid e_2 \mid e_4 \mid \cdots \mid e_{2\lfloor n/2 \rfloor}]^T$ is the $n \times n$ permutation matrix and e_i is the elementary column vector with the i th entry being 1 and the others entries being 0.

- (ii) *The matrix X is the adjacency matrix of the induced subgraph of $G_n(g, f)$ by the odd indexed vertex set $V_o = \{2i - 1 \mid 1 \leq i \leq \lceil n/2 \rceil\}$. In particular, the induced subgraph $\langle V_o \rangle$ is a Riordan graph of order $\lceil n/2 \rceil$ given by $G_{\lceil n/2 \rceil}(g'(\sqrt{z}), f(z))$.*
- (iii) *The matrix Y is the adjacency matrix of the induced subgraph of $G_n(g, f)$ by the even indexed vertex set $V_e = \{2i \mid 1 \leq i \leq \lfloor n/2 \rfloor\}$. In particular, the induced subgraph $\langle V_e \rangle$ is a Riordan graph of order $\lfloor n/2 \rfloor$ given by $G_{\lfloor n/2 \rfloor} \left(\left(\frac{gf}{z} \right)'(\sqrt{z}), f(z) \right)$.*
- (iv) *The matrix B representing the edges between V_o and V_e can be expressed as the sum of binary Riordan matrices as follows:*

$$B = \mathcal{B}(z \cdot (gf)'(\sqrt{z}), f(z))_{\lceil n/2 \rceil \times \lfloor n/2 \rfloor} + \mathcal{B}((zg)'(\sqrt{z}), f(z))_{\lfloor n/2 \rfloor \times \lceil n/2 \rceil}^T.$$

2.2 Families of Riordan graphs

There are many families of Riordan graphs introduced in [3]. Below we list those of them most relevant to this paper, along with examples of subfamilies.

Riordan graphs of the *Appell type*. This class of graphs is defined by an *Appell matrix* (g, z) , and thus it is precisely the class of Toeplitz graphs. Examples of graphs in this class are

- the null graphs N_n defined by $G_n(0, z)$;
- the path graphs P_n defined by $G_n(1, z)$;
- the complete graphs K_n defined by $G_n\left(\frac{1}{1-z}, z\right)$; and
- the complete bipartite graphs $K_{\lfloor \frac{n}{2} \rfloor, \lceil \frac{n}{2} \rceil}$ defined by $G_n\left(\frac{1}{1-z^2}, z\right)$.

Riordan graphs of the *Bell type*. This class of graphs is defined by a *Bell matrix* (g, zg) . Examples of graphs in this class are

- the null graphs N_n defined by $G_n(0, 0)$;
- the path graphs P_n defined by $G_n(1, z)$;

- the Pascal graphs PG_n defined by $G_n\left(\frac{1}{1-z}, \frac{z}{1-z}\right)$; and
- the Catalan graphs CG_n defined by $G_n\left(\frac{1-\sqrt{1-4z}}{2z}, \frac{1-\sqrt{1-4z}}{2}\right)$.

Riordan graphs of the checkerboard type. This class of graphs is defined by a checkerboard matrix (g, f) such that g is an even function and f is an odd function. Examples of graphs in this class are

- the null graphs N_n defined by $G_n(0, c)$ for any constant c ;
- the path graphs P_n defined by $G_n(1, z)$; and
- the complete bipartite graphs $K_{\lfloor \frac{n}{2} \rfloor, \lceil \frac{n}{2} \rceil}$ defined by $G_n\left(\frac{1}{1-z^2}, z\right)$.

Riordan graphs of the derivative type. This class of graphs is defined by functions (f', f) . Examples of graphs in this class are

- the null graphs N_n defined by $G_n(0, c)$ for any constant c ; and
- the path graphs P_n defined by $G_n(1, z)$.

o-decomposable Riordan graphs standing for *odd decomposable Riordan graphs*. This class of graphs is defined by requiring in (3) $Y = O$, where O is the zero matrix of Y 's size.

e-decomposable Riordan graphs standing for *even decomposable Riordan graphs*. This class of graphs is defined by requiring in (3) $X = O$.

2.3 Several known results on Riordan graphs

The following lemma is a corollary of Theorem 2.4.

Lemma 2.5 ([3]). *Every adjacency matrix $\mathcal{A} = \mathcal{A}_n(g, zg)$ of a Riordan graph $G_n(g, zg)$ of the Bell type can be expressed as the block matrix*

$$P\mathcal{A}P^T = \begin{bmatrix} X & B \\ B^T & O \end{bmatrix}, \quad (4)$$

where

$$X = \mathcal{A}_{\lceil n/2 \rceil}(g'(\sqrt{z}), zg(z))$$

and

$$B \equiv \mathcal{B}(zg(z), zg(z))_{\lceil n/2 \rceil, \lfloor n/2 \rfloor} + \mathcal{B}((zg)'(\sqrt{z}), zg(z))_{\lfloor n/2 \rfloor, \lceil n/2 \rceil}^T.$$

Thus, any Riordan graph of the Bell type is o-decomposable.

The following lemma is given by Theorem 3.6 (iv) in [3].

Lemma 2.6 ([3]). *A graph G_n is a Riordan graph of the checkerboard type if and only if G_n is bipartite with bipartitions V_o and V_e .*

In this paper, we also need the following lemmas.

Lemma 2.7 ([18]). *For the Pascal graph PG_n , we have $m(PG_{2^k}) = 3^k - 2^k$ and $m(PG_{2^k+1}) = 3^k$.*

Lemma 2.8 ([3]). *For the Catalan graph CG_n , we have $m(CG_{2^k}) = \frac{3^k-1}{2}$ and $m(CG_{2^k+1}) = \frac{3^k-1}{2} + 2^k$.*

Definition 2.9 ([3]). Let $G_n = G_n(g, f)$ be a proper Riordan graph with the odd and even vertex sets V_o and V_e , respectively.

- If $\langle V_o \rangle \cong G_{\lceil n/2 \rceil}(g, f)$ and $\langle V_e \rangle$ is a null graph then G_n is *io-decomposable*.

- If $\langle V_o \rangle$ is a null graph and $\langle V_e \rangle \cong G_{\lfloor n/2 \rfloor}(g, f)$ then G_n is *ie-decomposable*.

“io” and “ie” stand for “isomorphically odd” and “isomorphically even”, respectively.

Lemma 2.10 ([3]). *Let $G_n = G_n(g, zg)$ be an io-decomposable Riordan graph. Then, we have*

$$\chi(G_n) = \omega(G_n) = \lceil \log_2 n \rceil + 1,$$

where $\chi(G_n)$ and $\omega(G_n)$ are the chromatic number and clique number of G_n , respectively.

Lemma 2.11 ([3]). *A Riordan graph $G_n(g, zg)$ is io-decomposable if and only if*

$$g^2 \equiv g', \text{ i.e. } [z^j]g \equiv [z^{2j+1}]g.$$

2.4 The spectral graph theory

Let G be a graph and $\mathcal{A}(G)$ be its adjacency matrix. The eigenvalues $\lambda_1 \geq \dots \geq \lambda_n$ of $\mathcal{A}(G)$ are said to be the *eigenvalues* of G , and the eigenvalues form the *spectrum* of this graph. The largest eigenvalue λ_1 is called the *spectral radius* of G . The determinant of G , denoted by $\det(G)$, is the determinant of $\mathcal{A}(G)$. The number of positive and negative eigenvalues of a graph are called *positive* and *negative inertias* of the graph, respectively. We denote the positive and negative inertias of a Riordan graph G_n by $n^+(G_n)$ and $n^-(G_n)$, respectively. If G_n is understood from the context, we simply write n^+ and n^- for the inertias.

Let M be an $i \times j$ matrix. The *null space* of M is the set of all j -dimensional column vectors X such that $MX = 0$. The *dimension* of the null space of M is called the *nullity* of the matrix M , and is denoted by $\eta(M)$, or just by η when M is understood from the context. The nullity of G is $\eta(G) = \eta(\mathcal{A}(G))$. A graph G is called *singular* if its adjacency matrix is singular.

Definition 2.12. The *Laplacian matrix* $L(G)$ and *signless Laplacian matrix* $Q(G)$ of a graph G are defined, respectively, as $D(G) - \mathcal{A}(G)$ and $D(G) + \mathcal{A}(G)$, where $D(G)$ is the diagonal matrix of vertex degrees of G .

Definition 2.13. The *Laplacian spectrum* of a graph G is the sequence of its Laplacian eigenvalues, i.e. $\mu_1 \geq \mu_2 \geq \dots \geq \mu_{n-1} \geq \mu_n = 0$. We let $Lspec(G)$ denote the set $\{\mu_1, \mu_2, \dots, \mu_n\}$. The *signless Laplacian spectrum* of G is the sequence of its signless Laplacian eigenvalues, i.e. $q_1 \geq q_2 \geq \dots \geq q_n$.

It is well known that the average degree $\frac{2m}{n}$ of a graph G with n vertices and m edges is a lower bound for the spectral radius $\lambda_1(G)$, i.e.

$$\lambda_1(G) \geq \frac{2m}{n}. \quad (5)$$

Moreover, for the signless Laplacian spectral radius $q_1(G)$, we have

$$q_1(G) \geq 2\lambda(G) \geq \frac{4m}{n}. \quad (6)$$

The following lemma gives a relation between the eigenvalues of a real symmetric matrix and the eigenvalues of its *partitioned matrix*.

Lemma 2.14 ([5]). *Let A be a real symmetric matrix with eigenvalues $\lambda_1 \geq \lambda_2 \geq \dots \geq \lambda_n$. Given a partition $\{1, 2, \dots, n\} = V_1 \dot{\cup} V_2 \dot{\cup} \dots \dot{\cup} V_k$ with $|V_i| = n_i > 0$, consider the corresponding blocking $A = (A_{i,j})$, where $A_{i,j}$ is an $n_i \times n_j$ block and $1 \leq i, j \leq k$. Let $e_{i,j}$ be the sum of the entries in $A_{i,j}$ and set the matrix $B := (e_{i,j}/n_i)$ for $1 \leq i, j \leq k$. Then the eigenvalues of B interlace those of A , i.e. $\lambda_i \geq \rho_i \geq \lambda_{n-k-i}$ for $1 \leq i \leq k$, where ρ_i is the i th largest eigenvalue of B .*

The following lemma gives a relation between the adjacency matrix of a graph G and the *clique number* $\omega(G)$ of G , which is the size of a maximal clique in G .

Lemma 2.15 ([5]). *If G is a graph with the adjacency matrix A and the clique number $\omega(G)$, then*

$$\max \{X^T A X ; X \geq 0, \mathbf{1}^T X = 1\} = 1 - \frac{1}{\omega(G)},$$

where $\mathbf{1}$ is the vector of the proper size that contains only 1s .

In this paper, we also need the following lemmas.

Lemma 2.16 ([5]). *Let M be a symmetric $n \times n$ matrix with real entries. If $M = \begin{bmatrix} P & Q \\ Q^T & R \end{bmatrix}$, then $\lambda_1(M) + \lambda_n(M) \leq \lambda_1(P) + \lambda_1(R)$, where $\lambda_1(M)$ and $\lambda_n(M)$ stand for the largest and smallest eigenvalues of M , respectively.*

Lemma 2.17 ([1]). *Let A and B be real matrices of orders $n \times n$ and $n \times m$, respectively, and let $r = \text{rank}(B)$ and $\mathcal{A} := \begin{bmatrix} A & B \\ B^T & O \end{bmatrix}$. Then $n^-(\mathcal{A}) \geq r$, $\eta(\mathcal{A}) \geq 0$ and $n^+(\mathcal{A}) \geq r$. If, in addition, $n = m$ and B is nonsingular, then $n^-(\mathcal{A}) = n^+(\mathcal{A}) = n$ and $\eta(\mathcal{A}) = 0$.*

Lemma 2.18 ([1]). *Let $A \in \mathbb{R}^{n \times n}$, $B \in \mathbb{R}^{m \times m}$, $C \in \mathbb{R}^{n \times m}$, and A and B be symmetric. Also, let $\mathcal{A}_0 := \begin{bmatrix} A & O \\ O & B \end{bmatrix}$, $\mathcal{A} := \begin{bmatrix} A & C \\ C^T & B \end{bmatrix}$, $\sigma_{\max}(C)$ be the largest singular value of C , and $\eta := \min_{\substack{i=1, \dots, n \\ j=1, \dots, m}} |\lambda_i(A) - \lambda_j(B)|$. Then, for all $i = 1, \dots, n + m$,*

$$|\lambda_i(\mathcal{A}) - \lambda_i(\mathcal{A}_0)| \leq \frac{2\sigma_{\max}^2(C)}{\eta + \sqrt{\eta^2 + 4\sigma_{\max}^2(C)}}.$$

Lemma 2.19 ([5, 19]). *For any graph G with chromatic number $\chi(G)$ and clique number $\omega(G)$, we have*

- (i) $\chi(G) - 1 \leq \lambda_1(G) \leq n \left(1 - \frac{1}{\omega(G)}\right)$.
- (ii) $\chi(G) \geq 1 + \frac{\lambda_1(G)}{|\lambda_n(G)|}$.
- (iii) $\chi(G) \geq 1 + \frac{\lambda_1(G)}{\mu_1(G) - \lambda_1(G)}$.

Lemma 2.20 ([14]). *Let G be a graph of order n . Then $\mu_1(G) \leq n$, where the equality holds if and only if the complement \overline{G} of G is disconnected. Moreover, $L\text{spec}(\overline{G}) = \{n - \mu_{n-1}, n - \mu_{n-2}, \dots, n - \mu_1, 0\}$.*

Lemma 2.21 ([11]). *Let G ($\not\cong K_n$) be a graph of order n . Then $\mu_{n-1}(G) \leq \delta(G)$, where $\delta(G)$ is the minimum vertex degree in G .*

Lemma 2.22 ([14]). *Let G be a graph on n vertices with at least one edge. Then $\mu_1(G) \geq \Delta(G) + 1$, where $\Delta(G)$ is the maximum vertex degree in G . Moreover, if G is connected, then the equality holds if and only if $\Delta(G) = n - 1$.*

The following result states for which n the Catalan number C_n is odd.

Lemma 2.23 ([10]). *The only Catalan numbers C_n that are odd are those for which $n = 2^k - 1$ for $k \geq 0$. In particular, $C_{2i} \equiv 0$ for $i \in \mathbb{N}$.*

3 Eigenvalues of Riordan graphs

In this section we present several lower and upper bounds on the eigenvalues and the (signless) Laplacian eigenvalues of Riordan graphs, which will improve (5) and (6) for this class of graphs. To achieve our goals, we begin with a general result valid for all graphs.

Theorem 3.1. *Let $G = (V, E)$ be a graph with $|V| = n$ and $|E| = m$, and let W be a nonempty subset of V with $|W| = k$. Then*

$$\begin{aligned}
\text{(i)} \quad \lambda_1(G) &\geq \frac{m_1}{k} + \frac{m_2}{n-k} + \sqrt{\left(\frac{m_1}{k} - \frac{m_2}{n-k}\right)^2 + \frac{(m - m_1 - m_2)^2}{k(n-k)}}. \\
\text{(ii)} \quad \mu_1(G) &\geq \frac{n(m - m_1 - m_2)}{k(n-k)}. \\
\text{(iii)} \quad q_1(G) &\geq \frac{1}{2} \left(c_1 + c_2 + \sqrt{(c_1 - c_2)^2 + \frac{4(m - m_1 - m_2)^2}{k(n-k)}} \right). \\
\text{(iv)} \quad q_2(G) &\geq \frac{1}{2} \left(c_1 + c_2 - \sqrt{(c_1 - c_2)^2 + \frac{4(m - m_1 - m_2)^2}{k(n-k)}} \right) \geq q_n(G)
\end{aligned}$$

where $c_1 = \frac{m+3m_1-m_2}{k}$, $c_2 = \frac{m-m_1+3m_2}{n-k}$, $m_1 = m(\langle W \rangle)$ and $m_2 = m(\langle V \setminus W \rangle)$.

Proof: Let \mathcal{W} and \mathcal{U} be, respectively, the adjacency matrices of the induced subgraphs $\langle W \rangle$ and $\langle V \setminus W \rangle$. Let \mathcal{A} be the adjacency matrix of G . Then there exists a permutation matrix P such that

$$PAP^T = \begin{bmatrix} \mathcal{W} & \mathcal{F} \\ \mathcal{F}^T & \mathcal{U} \end{bmatrix}.$$

Using Lemma 2.14 on PAP^T , we obtain

$$B_1 := \begin{bmatrix} \frac{2m_1}{k_1} & \frac{m_3}{k_1} \\ \frac{m_3}{k_2} & \frac{2m_2}{k_2} \end{bmatrix}$$

where $k_1 := |W|$ and $k_2 := n - k_1 = |V \setminus W|$, and m_3 stands for the number of 1s in the submatrix \mathcal{F} , which is the number of edges between W and $V \setminus W$. It is not difficult to see that the eigenvalues of B_1 are given by

$$\lambda_1(B_1) = \frac{m_1}{k_1} + \frac{m_2}{k_2} + \sqrt{\left(\frac{m_1}{k_1} - \frac{m_2}{k_2}\right)^2 + \frac{m_3^2}{k_1 k_2}}$$

and

$$\lambda_2(B_1) = \frac{m_1}{k_1} + \frac{m_2}{k_2} - \sqrt{\left(\frac{m_1}{k_1} - \frac{m_2}{k_2}\right)^2 + \frac{m_3^2}{k_1 k_2}}.$$

By Lemma 2.14, we have $\lambda_1(G) \geq \lambda_1(B_1)$. This completes the proof of (i).

Suppose that $L = D - \mathcal{A}$ is the Laplacian matrix of G . Then by the same permutation matrix P as above, we obtain

$$PLP^T = PDP^T - PAP^T = \begin{bmatrix} D_{\mathcal{W}} & O \\ O^T & D_{\mathcal{U}} \end{bmatrix} - \begin{bmatrix} \mathcal{W} & \mathcal{F} \\ \mathcal{F}^T & \mathcal{U} \end{bmatrix} = \begin{bmatrix} D_{\mathcal{W}} - \mathcal{W} & -\mathcal{F} \\ -\mathcal{F}^T & D_{\mathcal{U}} - \mathcal{U} \end{bmatrix}$$

where $D_{\mathcal{W}}$ and $D_{\mathcal{U}}$ are, respectively, the degree matrices with the same row indices as \mathcal{W} and \mathcal{U} , and O is the zero matrix of the proper size. Again, using Lemma 2.14 on PLP^T , we obtain

$$B_2 := \begin{bmatrix} \frac{m_3}{k_1} & -\frac{m_3}{k_1} \\ -\frac{m_3}{k_2} & \frac{m_3}{k_2} \end{bmatrix}.$$

It is easy to see that the eigenvalues of B_2 are given by

$$\lambda_1(B_2) = \frac{nm_3}{k_1 k_2} \quad \text{and} \quad \lambda_2(B_2) = 0.$$

By Lemma 2.14, we have $\mu_1(G) \geq \lambda_1(B_2)$. Hence, (ii) is proved.

Suppose that $Q = D + A$ is the signless Laplacian matrix of G . Then, by the same permutation matrix P as above, we obtain

$$PQP^T = PDP^T + PAP^T = \begin{bmatrix} D_{\mathcal{W}} & O \\ O^T & D_{\mathcal{U}} \end{bmatrix} + \begin{bmatrix} \mathcal{W} & \mathcal{F} \\ \mathcal{F}^T & \mathcal{U} \end{bmatrix} = \begin{bmatrix} D_{\mathcal{W}} + \mathcal{W} & \mathcal{F} \\ \mathcal{F}^T & D_{\mathcal{U}} + \mathcal{U} \end{bmatrix}.$$

Using Lemma 2.14 on PQP^T , we obtain

$$B_3 = \begin{bmatrix} \frac{4m_1+m_3}{k_1} & \frac{m_3}{k_1} \\ \frac{m_3}{k_2} & \frac{4m_2+m_3}{k_2} \end{bmatrix}.$$

The eigenvalues of B_3 are given by

$$\lambda_1(B_3) = \frac{1}{2} \left(\frac{4m_1+m_3}{k_1} + \frac{4m_2+m_3}{k_2} + \sqrt{\left(\frac{4m_1+m_3}{k_1} - \frac{4m_2+m_3}{k_2} \right)^2 + \frac{4m_3^2}{k_1k_2}} \right)$$

and

$$\lambda_2(B_3) = \frac{1}{2} \left(\frac{4m_1+m_3}{k_1} + \frac{4m_2+m_3}{k_2} - \sqrt{\left(\frac{4m_1+m_3}{k_1} - \frac{4m_2+m_3}{k_2} \right)^2 + \frac{4m_3^2}{k_1k_2}} \right).$$

One can see that $\lambda_2(B_3) > 0$. By Lemma 2.14, we have $q_1(G) \geq \lambda_1(B_3)$ and $q_2(G) \geq \lambda_2(B_3) \geq q_n(G)$. This completes the proofs for (iii) and (iv). \square

Corollary 3.2. *Let $G = (V, E)$ be a graph with $|V| = n$ and $|E| = m$, and let W be a nonempty subset of V with $|W| = k$. Also, let $m_1 = m(\langle W \rangle)$ and $m_2 = m(\langle V \setminus W \rangle)$, and $m > m_1 + m_2 + 2\sqrt{m_1 m_2}$. Then $\lambda_n(G) < 0$.*

Proof: For any graph G , $\lambda_2(B_1) \geq \lambda_n(G)$, where B_1 and $\lambda_2(B_1)$ can be found in the proof of Theorem 3.1. Since $m > m_1 + m_2 + 2\sqrt{m_1 m_2}$, we have

$$\lambda_n(G) \leq \frac{m_1}{k} + \frac{m_2}{n-k} - \sqrt{\left(\frac{m_1}{k} - \frac{m_2}{n-k} \right)^2 + \frac{(m - m_1 - m_2)^2}{k(n-k)}} < 0.$$

\square

The following theorem, the main result in this section, follows from Theorems 2.4 and 3.1.

Theorem 3.3. *Let $G_n = G_n(g, f)$ be a Riordan graph with n vertices and m edges. Then*

$$(i) \quad \lambda_1(G_n) \geq \frac{m_1}{n_1} + \frac{m_2}{n-n_1} + \sqrt{\left(\frac{m_1}{n_1} - \frac{m_2}{n-n_1} \right)^2 + \frac{(m - m_1 - m_2)^2}{n_1(n-n_1)}}. \quad (7)$$

$$(ii) \quad \mu_1(G_n) \geq \frac{n(m - m_1 - m_2)}{n_1(n - n_1)}.$$

$$(iii) \quad q_1(G_n) \geq \frac{1}{2} \left(c_1 + c_2 + \sqrt{(c_1 - c_2)^2 + \frac{4(m - m_1 - m_2)^2}{n_1(n - n_1)}} \right). \quad (8)$$

$$(iv) \quad q_2(G_n) \geq \frac{1}{2} \left(c_1 + c_2 - \sqrt{(c_1 - c_2)^2 + \frac{4(m - m_1 - m_2)^2}{n_1(n - n_1)}} \right) \geq q_n(G),$$

where $n_1 = \lceil \frac{n}{2} \rceil$, $m_1 = m(G_{n_1}(g'(\sqrt{z}), f))$, $m_2 = m(G_{n-n_1}((\frac{gf}{z})'(\sqrt{z}), f))$, $c_1 = \frac{m+3m_1-m_2}{n_1}$ and $c_2 = \frac{m-m_1+3m_2}{n-n_1}$.

Proposition 3.4. For a Riordan graph G_n , the lower bound in (7) is greater than or equal to the lower bound in (5).

Proof: Suppose that $n = 2k$, and m_3 is defined as in the proof of Theorem 3.1. We have

$$\begin{aligned}
(m_1 - m_2)^2 \geq 0 &\Rightarrow (m_1 + m_2)^2 \geq 4m_1m_2 \Rightarrow \\
m_3^2 &\geq m_3^2 + (m_1 + m_2)^2 + 2m_3(m_1 + m_2) + 4m_1m_2 - 2(m_1 + m_2)^2 - 2m_3(m_1 + m_2) \\
&\Rightarrow m_3^2 \geq m^2 + 4m_1m_2 - 2mm_1 - 2mm_2 \Rightarrow \\
\frac{(m_1 - m_2)^2 + m_3^2}{k^2} &\geq \frac{m^2 + (m_1 + m_2)^2 - 2m(m_1 + m_2)}{k^2} \Rightarrow \\
\frac{m_1 + m_2}{k} + \sqrt{\frac{(m_1 - m_2)^2 + m_3^2}{k^2}} &\geq \frac{m}{k} \Rightarrow \\
\frac{m_1}{n_1} + \frac{m_2}{n_2} + \sqrt{\left(\frac{m_1}{n_1} - \frac{m_2}{n_2}\right)^2 + \frac{m_3^2}{n_1n_2}} &\geq \frac{2m}{n}
\end{aligned}$$

as $n_1 = n_2 = k$. Similarly, the result for $n = 2k + 1$ follows. \square

Proposition 3.5. For a Riordan graph G_n , the lower bound in (8) is greater than or equal to the lower bound in (6).

Proof: Since $q_1 \geq 2\lambda_1$ for any graph, the result follows from Proposition 3.4. \square

The following results are obtained by Theorems 2.4 and 3.3.

Corollary 3.6. Let $G_n = G_n(g, z)$ be a Riordan graph of the Appell type with n vertices and m edges. Let n be even and $m_1 = m(G_{n/2}(g'(\sqrt{z}), z))$. Then

- (i) The lower bounds in (7) and (5) coincide for the spectral radius $\lambda_1(G_n)$.
- (ii) The lower bounds in (8) and (6) coincide for the signless Laplacian spectral radius $q_1(G_n)$.
- (iii) $\mu_1(G_n) \geq \frac{4(m-2m_1)}{n}$.
- (iv) $q_2(G_n) \geq \frac{8m_1}{n} \geq q_n(G)$.
- (v) If $m > 4m_1$, then $\lambda_n(G_n) \leq \frac{8m_1-2m}{n} < 0$.

Corollary 3.7. Let $G_n = G_n(g, f)$ be a Riordan graph with n vertices and m edges. If $[z^{2j+1}]g \equiv 0$ for all $j \geq 0$, then

$$\begin{aligned}
\text{(i)} \quad \lambda_1(G_n) &\geq \frac{m_2}{n - n_1} + \sqrt{\frac{m_2^2}{(n - n_1)^2} + \frac{(m - m_2)^2}{n_1(n - n_1)}} \\
\text{(ii)} \quad \mu_1(G_n) &\geq \frac{n(m - m_2)}{n_1(n - n_1)} \\
\text{(iii)} \quad q_1(G_n) &\geq \frac{1}{2} \left(c_1 + c_2 + \sqrt{(c_1 - c_2)^2 + \frac{4(m - m_2)^2}{n_1(n - n_1)}} \right) \\
\text{(iv)} \quad q_2(G_n) &\geq \frac{1}{2} \left(c_1 + c_2 - \sqrt{(c_1 - c_2)^2 + \frac{4(m - m_2)^2}{n_1(n - n_1)}} \right) \geq q_n(G)
\end{aligned}$$

where $n_1 = \lceil \frac{n}{2} \rceil$, $m_2 = m(G_{n-n_1}(\frac{gf}{z})(\sqrt{z}), f)$, $c_1 = \frac{m-m_2}{n_1}$ and $c_2 = \frac{m+3m_2}{n-n_1}$.

The following result is obtained by Lemma 2.6 and Corollary 3.7.

Corollary 3.8. *Let $G_n = G_n(g, f)$ be a Riordan graph of the checkerboard type with n vertices and m edges, and let $n_1 = \lceil \frac{n}{2} \rceil$. Then,*

- (i) $\lambda_1(G_n) = -\lambda_n(G_n) \geq \frac{m}{\sqrt{n_1(n-n_1)}}$.
- (ii) $\mu_1(G_n) = q_1(G_n) \geq \frac{nm}{n_1(n-n_1)}$.
- (iii) $q_n(G_n) = 0$.

Corollary 3.9. *Let $G_n = G_n(g, f)$ be a Riordan graph with n vertices and m edges such that $[z^{2j}]g = [z^{2j}]f \equiv 0$ for all $j \geq 0$. Then G_n is disconnected with components $H_1 \cong G_{n_1}(g'(\sqrt{z}), f)$ and $H_2 \cong G_{n-n_1}((\frac{gf}{z})'(\sqrt{z}), f)$, where $n_1 = \lceil \frac{n}{2} \rceil$. Moreover, letting $m_1 = m(H_1)$, we have*

- (i) $\lambda_1(G_n) = \max\{\lambda_1(H_1), \lambda_1(H_2)\} \geq \max\left\{\frac{2m_1}{n_1}, \frac{2(m-m_1)}{n-n_1}\right\}$.
- (ii) $q_1(G_n) = \max\{q_1(H_1), q_1(H_2)\} \geq \max\left\{\frac{4m_1}{n_1}, \frac{4(m-m_1)}{n-n_1}\right\}$.

The following result is obtained by Lemma 2.5 and Theorem 3.3.

Corollary 3.10. *Let $G_n = G_n(g(z), zg(z))$ be a Riordan graph of the Bell type with n vertices and m edges. Then,*

- (i) $\lambda_1(G_n) \geq \frac{m_1}{n_1} + \sqrt{\frac{m_1^2}{n_1^2} + \frac{(m-m_1)^2}{n_1(n-n_1)}}$
- (ii) $\mu_1(G_n) \geq \frac{n(m-m_1)}{n_1(n-n_1)}$
- (iii) $q_1(G_n) \geq \frac{1}{2} \left(c_1 + c_2 + \sqrt{(c_1 - c_2)^2 + \frac{4(m-m_1)^2}{n_1(n-n_1)}} \right)$
- (iv) $q_2(G_n) \geq \frac{1}{2} \left(c_1 + c_2 - \sqrt{(c_1 - c_2)^2 + \frac{4(m-m_1)^2}{n_1(n-n_1)}} \right) \geq q_n(G)$

where $n_1 = \lceil \frac{n}{2} \rceil$, $c_1 = \frac{m+3m_1}{n_1}$, $c_2 = \frac{m-m_1}{n-n_1}$ and

$$m_1 = \begin{cases} m(G_{n_1}(g'(\sqrt{z}), zg(z))) & \text{in the general case;} \\ m(G_{n_1}(g(z), zg(z))) & \text{if } G_n \text{ is io-decomposable.} \end{cases}$$

In the following proposition, we use Lemmas 2.7 and 2.8, and Corollary 3.10 to present lower bounds on some eigenvalues and (signless) Laplacian eigenvalues of PG_{2^k} , PG_{2^k+1} , CG_{2^k} and CG_{2^k+1} .

Proposition 3.11. *For $k \geq 2$, we have*

- (i) $\lambda_1(PG_{2^k}) \geq \frac{3^{k-1} + \sqrt{(3^{k-1} - 2^{k-1})^2 + 4(3^{k-1} - 2^{k-2})^2}}{2^{k-1}} - 1$.
- (ii) $q_1(PG_{2^k}) \geq \frac{1}{2} \left(\frac{3^{k-1}}{2^{k-4}} + \frac{\sqrt{(3^{k-1} - 2^{k-1})^2 + (3^{k-1} - 2^{k-2})^2}}{2^{k-3}} \right) - 3$.
- (iii) $\lambda_1(PG_{2^k+1}) \geq \frac{3^{k-1} \left(1 + \sqrt{17 + 2^{3-k}} \right)}{2^{k-1} + 1}$.

$$\begin{aligned}
\text{(iv)} \quad q_1(PG_{2^{k+1}}) &\geq \frac{3^{k-1} \left(2^{k+1} + 1 + \sqrt{2^{2k+1} + 1} \right)}{2^{k-1} (2^{k-1} + 1)}. \\
\text{(v)} \quad \lambda_1(CG_{2^k}) &\geq \frac{5 \cdot 3^{2k-2} - 3^{k-1}}{2^k}. \\
\text{(vi)} \quad \mu_1(CG_{2^k}) &\geq \frac{3^{k-1}}{2^{k-2}}. \\
\text{(vii)} \quad q_1(CG_{2^k}) &\geq \frac{2 \cdot 3^{k-1} - 1 + \sqrt{2 \cdot 3^{k-1} (3^{k-1} - 1) + 1}}{2^{k-1}}. \\
\text{(viii)} \quad q_2(CG_{2^k}) &\geq \frac{2 \cdot 3^{k-1} - 1 - \sqrt{2 \cdot 3^{k-1} (3^{k-1} - 1) + 1}}{2^{k-1}} \geq q_n(CG_{2^k}). \\
\text{(ix)} \quad \lambda_1(CG_{2^{k+1}}) &\geq \frac{b + \sqrt{b^2 + (1 + 2^{1-k})(3^{k-1} + 2^{k-1})^2}}{2^{k-1} + 1}, \text{ where } b = \frac{3^{k-1} + 2^{k-1}}{2}. \\
\text{(x)} \quad \mu_1(CG_{2^{k+1}}) &\geq \frac{(2^k + 1)(3^{k-1} + 2^{k-1})}{2^{2k-2} + 2^{k-1}}.
\end{aligned}$$

The following is a result on the spectral radius and the (signless) Laplacian spectral radius of an io-decomposable Riordan graph $G_n = G_n(g, zg)$.

Theorem 3.12. *Let $G_n = G_n(g, zg)$ be io-decomposable with n vertices and m edges. Then*

$$\begin{aligned}
\text{(i)} \quad m &\leq \frac{n^2}{2} \left(1 - \frac{1}{\lceil \log_2 n \rceil + 1} \right). \\
\text{(ii)} \quad \lceil \log_2 n \rceil &\leq \lambda_1(G_n) \leq n \left(1 - \frac{1}{\lceil \log_2 n \rceil + 1} \right). \\
\text{(iii)} \quad \frac{\lambda_1(G_n)}{|\lambda_n(G_n)|} &\leq \lceil \log_2 n \rceil. \\
\text{(iv)} \quad \lceil \log_2 n \rceil + 1 &\leq \frac{\lceil \log_2 n \rceil + 1}{\lceil \log_2 n \rceil} \lambda_1(G_n) \leq \mu_1(G_n). \\
\text{(v)} \quad \lambda_1(G_n) - \lambda_1(G_{\lceil \frac{n}{2} \rceil}) &\leq -\lambda_n(G_n). \\
\text{(vi)} \quad q_1(G_n) &\geq 2 \lceil \log_2 n \rceil.
\end{aligned}$$

Proof: Considering $X^T = \underbrace{[1/n, 1/n, \dots, 1/n]}_n$ in Lemma 2.15, and applying Lemma 2.10, we obtain

$$\frac{2}{n^2} \sum_{ij \in E(G_n)} 1 = \frac{2m}{n^2} \leq 1 - \frac{1}{\lceil \log_2 n \rceil + 1} \Rightarrow m \leq \frac{n^2}{2} \left(1 - \frac{1}{\lceil \log_2 n \rceil + 1} \right).$$

Hence, the proof of (i) is complete. We obtain (ii), (iii) and (iv) directly from Lemmas 2.10 and 2.19.

Let \mathcal{A}_n be the adjacency matrix of G_n . By Lemma 2.5 and the definition of an io-decomposable Riordan graph, we have $P^T \mathcal{A}_n P = \begin{bmatrix} \mathcal{A}_{\lceil \frac{n}{2} \rceil} & B \\ B^T & O \end{bmatrix}$. From this, after applying Lemma 2.16, we obtain $\lambda_1(\mathcal{A}_n) + \lambda_n(\mathcal{A}_n) \leq \lambda_1(\mathcal{A}_{\lceil \frac{n}{2} \rceil})$, which gives the desired result in (v). Finally, (vi) follows from (ii) and (6). \square

In what follows, we give an upper bound for the absolute value of the smallest eigenvalue of an io-decomposable Riordan graph G_n .

Theorem 3.13. Let $G_n = G_n(g, zg)$ be an io-decomposable Riordan graph with the adjacency matrix $\mathcal{A}_n := P \begin{bmatrix} \mathcal{A}_{\lceil \frac{n}{2} \rceil} & B \\ B^T & O \end{bmatrix} P^T$, and let $\lambda_i(G_n)$ and $\sigma_{\max}(B)$ be, respectively, the i th largest eigenvalue of G_n and the largest singular value of B . Also, let $\eta := \min_{i=1, \dots, \lceil \frac{n}{2} \rceil} |\lambda_i(G_{\lceil \frac{n}{2} \rceil})|$. Then

$$|\lambda_n(G_n)| \leq \frac{2\sigma_{\max}^2(B)}{\eta + \sqrt{\eta^2 + 4\sigma_{\max}^2(B)}}.$$

Proof: By Theorem 3.12 (v) and Lemma 2.18, we obtain

$$|\lambda_n(G_n)| \leq \max_{\substack{i=1, \dots, \lceil \frac{n}{2} \rceil \\ j=1+\lceil \frac{n}{2} \rceil, \dots, n}} \left\{ |\lambda_i(G_n) - \lambda_i(G_{\lceil \frac{n}{2} \rceil})|, |\lambda_j(G_n)| \right\} \leq \frac{2\sigma_{\max}^2(B)}{\eta + \sqrt{\eta^2 + 4\sigma_{\max}^2(B)}}.$$

□

The following result is related to the structure of the adjacency matrix of an io-decomposable Riordan graph $G_n(g, zg)$, and it follows from the definition of an io-decomposable Riordan graph of the Bell type and Lemma 2.5.

Lemma 3.14. Let $\mathcal{A}_n := P \begin{bmatrix} \mathcal{A}_{\lceil \frac{n}{2} \rceil} & B \\ B^T & O \end{bmatrix} P^T$ be the adjacency matrix of an io-decomposable Riordan graph $G_n(g, zg)$, and let $b_{i,j}$ be the (i, j) th entry of B^T . If n is odd, then $P^T \mathcal{A}_n P$ is given by

$$\begin{bmatrix} 0 & r_{1,2} & \cdots & r_{1,k} & b_{1,1} & \cdots & b_{k-1,1} \\ r_{2,1} & 0 & \ddots & \vdots & r_{2,1} & \ddots & \vdots \\ \vdots & \ddots & \ddots & \vdots & \vdots & \ddots & \vdots \\ r_{k,1} & \cdots & r_{k,k-1} & 0 & r_{k,1} & \cdots & r_{k,k-1} \\ \hline b_{1,1} & r_{1,2} & \cdots & r_{1,k} & 0 & \cdots & 0 \\ \vdots & \ddots & \ddots & \vdots & \vdots & \ddots & \vdots \\ b_{k-1,1} & \cdots & b_{k-1,k-1} & r_{k-1,k} & 0 & \cdots & 0 \end{bmatrix}$$

where $k = \frac{n+1}{2}$. Otherwise, if n is even and $j = \frac{n}{2}$, then $P^T \mathcal{A}_n P$ is

$$\begin{bmatrix} 0 & r_{1,2} & \cdots & r_{1,j} & b_{1,1} & \cdots & \cdots & b_{j,1} \\ r_{2,1} & 0 & \ddots & \vdots & r_{2,1} & \ddots & \vdots & \vdots \\ \vdots & \ddots & \ddots & \vdots & \vdots & \ddots & \ddots & \vdots \\ r_{j,1} & \cdots & r_{j,j-1} & 0 & r_{j,1} & \cdots & r_{j,j-1} & b_{j,j} \\ \hline b_{1,1} & r_{1,2} & \cdots & r_{1,j} & 0 & \cdots & \cdots & 0 \\ \vdots & \ddots & \ddots & \vdots & \vdots & \ddots & \vdots & \vdots \\ \vdots & \vdots & \ddots & \vdots & \vdots & \ddots & \ddots & \vdots \\ b_{j,1} & \cdots & \cdots & b_{j,j} & 0 & \cdots & \cdots & 0 \end{bmatrix}.$$

In the following, we give a lower bound on the spectral radius of any io-decomposable Riordan graph $G_n(g, zg)$. We shall mean

$$\{gf^k\}_n(1) := \sum_{j=k}^n [z^j] \{gf^k\}$$

by the substitution of $z = 1$ in the Taylor expansion in z of $\{gf^k\}$ up to degree n modulo 2.

Theorem 3.15. *Let $G_n = G_n(g, zg)$ be an io-decomposable Riordan graph. Then,*

$$\lambda_1(G_n) \geq \frac{1 + \sqrt{2}}{n_1} \sum_{j=0}^{n_1-2} \{g'(\sqrt{z}) \cdot f^j\}_{n_1-2}(1).$$

Proof: By Corollary 3.10, we have

$$\lambda_1(G_n) \geq \frac{m_1}{n_1} + \sqrt{\frac{m_1^2}{n_1^2} + \frac{(m - m_1)^2}{n_1(n - n_2)}}. \quad (9)$$

Let $\mathcal{A}_n = P \begin{bmatrix} \mathcal{A}_{\lceil \frac{n}{2} \rceil} & B \\ B^T & \mathcal{O} \end{bmatrix} P^T$ be the adjacency matrix of G_n . Then, by Lemma 3.14, we have

$$\sigma(B) > m \left(\mathcal{A}_{\lceil \frac{n}{2} \rceil} \right) = m_1 \geq \sqrt{\frac{n - n_1}{n_1}} m_1 \Rightarrow \sigma(B)^2 > \frac{n - n_1}{n_1} m_1^2 \Rightarrow \frac{(m - m_1)^2}{n_1(n - n_1)} > \frac{m_1^2}{n_1^2}.$$

From (9), the above and Lemma 2.5, the required result is obtained. \square

The following two results give lower bounds on the largest Laplacian eigenvalue $\mu_1(G_n)$ of a Riordan graph G_n .

Theorem 3.16. *Let $G_n = G_n(g, f)$ be a Riordan graph with n vertices and m edges, and let $n_1 = \lceil \frac{n}{2} \rceil$ and $n_2 = n - n_1$. Then*

$$\mu_1(G_n) \geq \frac{n}{n_1 n_2} \left(\sum_{j=0}^{n_2-1} \{z(gf)'(\sqrt{z}) \cdot f^j\}_{n_1-1}(1) + \sum_{j=0}^{n_1-1} \{(zg)'(\sqrt{z}) \cdot f^j\}_{n_2-1}(1) \right).$$

Proof: Let $L = D - \mathcal{A}$ be the Laplacian matrix of G_n . Using the same technique as in the proof of Theorem 3.1, and by Lemma 2.14 applied to PLP^T , we obtain

$$B_2 := \begin{bmatrix} \frac{\sigma(B)}{n_1} & -\frac{\sigma(B)}{n_1} \\ -\frac{\sigma(B)}{n_2} & \frac{\sigma(B)}{n_2} \end{bmatrix},$$

where $\sigma(B)$ denotes the number of 1s in the matrix B. It is easy to see that the eigenvalues of B_2 are given by

$$\lambda_1(B_2) = \frac{n\sigma(B)}{n_1 n_2} \quad \text{and} \quad \lambda_2(B_2) = 0.$$

By Lemma 2.14, we have

$$\mu_1(G) \geq \lambda_1(B_2) = \frac{n\sigma(B)}{n_1 n_2}. \quad (10)$$

Now, by (iv) in the Riordan Graph Decomposition Theorem (Theorem 2.4), we have

$$\begin{aligned} \sigma(B) &= \sigma(\mathcal{B}(z \cdot (gf)'(\sqrt{z}), f(z))_{n_1 \times n_2}) + \sigma(\mathcal{B}((zg)'(\sqrt{z}), f(z))_{n_2 \times n_1}^T) \\ &= \sigma(\mathcal{B}(z \cdot (gf)'(\sqrt{z}), f(z))_{n_1 \times n_2}) + \sigma(\mathcal{B}((zg)'(\sqrt{z}), f(z))_{n_2 \times n_1}) \\ &= \sum_{j=0}^{n_2-1} \{z(gf)'(\sqrt{z}) \cdot f^j\}_{n_1-1}(1) + \sum_{j=0}^{n_1-1} \{(zg)'(\sqrt{z}) \cdot f^j\}_{n_2-1}(1). \end{aligned}$$

Using this in (10), the required result is obtained. \square

The following result is an immediate corollary of Theorem 3.16.

Corollary 3.17. *Let $G_n = G_n(g, f)$ be a proper Riordan graph with n vertices and m edges. Then*

$$\mu_1(G_n) \geq \frac{n}{n_1 n_2} \left(\{z(gf)'(\sqrt{z})\}_{n_1-1} (1) + \{(zg)'(\sqrt{z})\}_{n_2-1} (1) + 2 \left\lfloor \frac{n}{2} \right\rfloor - 3 \right).$$

For a graph G , let $a(G)$ be the *algebraic connectivity* of G . The following result gives a relation between the algebraic connectivity of $\langle V_o \rangle$ and the median Laplacian eigenvalues $\mu_{\lceil \frac{n}{2} \rceil}(G_n)$ and $\mu_{\lceil \frac{n}{2} \rceil+1}(G_n)$ of an o -decomposable Riordan graph G_n .

Theorem 3.18. *Let $G_n = G_n(g, f)$ be an o -decomposable Riordan graph and let $H \cong \langle V_o \rangle$. Then*

$$\mu_{\lceil \frac{n}{2} \rceil+1}(G_n) \leq \left\lfloor \frac{n}{2} \right\rfloor \quad (11)$$

$$\mu_{\lceil \frac{n}{2} \rceil}(G_n) \leq \begin{cases} \left\lfloor \frac{n}{2} \right\rfloor + a(H) & \text{if } n \text{ is even or } n \text{ is odd with } a(H) > 1; \\ \left\lceil \frac{n}{2} \right\rceil & \text{if } n \text{ is odd with } a(H) \leq 1. \end{cases} \quad (12)$$

Proof: One can see that $K_{\lfloor \frac{n}{2} \rfloor}$ is a subgraph of \overline{G}_n . Further, it is well known [5] that the Laplacian spectrum of $K_{\lfloor \frac{n}{2} \rfloor}$ is as follows:

$$L\text{spec} \left(K_{\lfloor \frac{n}{2} \rfloor} \right) = \left\{ \underbrace{\left\lfloor \frac{n}{2} \right\rfloor, \dots, \left\lfloor \frac{n}{2} \right\rfloor}_{\lfloor \frac{n}{2} \rfloor - 1}, 0 \right\}.$$

From the above, we obtain

$$\mu_{\lfloor \frac{n}{2} \rfloor - 1}(\overline{G}_n) \geq \mu_{\lfloor \frac{n}{2} \rfloor - 1} \left(K_{\lfloor \frac{n}{2} \rfloor} \right) = \left\lfloor \frac{n}{2} \right\rfloor.$$

From this and Lemma 2.20, we obtain

$$n - \mu_{\lceil \frac{n}{2} \rceil+1}(G_n) \geq \left\lfloor \frac{n}{2} \right\rfloor,$$

which completes the proof of (11).

Now, we consider the subgraph $\overline{H} \cup K_{\lfloor \frac{n}{2} \rfloor}$ of \overline{G}_n , where $H \cong \langle V_o \rangle$. By Lemma 2.20, we have $\mu_i(\overline{H}) = \left\lceil \frac{n}{2} \right\rceil - \mu_{\lceil \frac{n}{2} \rceil - i}(H)$ for $1 \leq i \leq \left\lceil \frac{n}{2} \right\rceil - 1$. From this and Lemma 2.20, we have

$$\mu_{\lfloor \frac{n}{2} \rfloor}(\overline{G}_n) \geq \min \left\{ \left\lfloor \frac{n}{2} \right\rfloor, \mu_1(\overline{H}) \right\} = \min \left\{ \left\lfloor \frac{n}{2} \right\rfloor, \left\lceil \frac{n}{2} \right\rceil - a(H) \right\}. \quad (13)$$

If n is even, from (13) and the fact that $\mu_1(\overline{H}) \leq |V(\overline{H})| = \frac{n}{2}$, we obtain

$$\mu_{\frac{n}{2}}(\overline{G}_n) \geq \mu_1(\overline{H}) \Rightarrow n - \mu_{\frac{n}{2}}(G_n) \geq \frac{n}{2} - \mu_{\frac{n}{2}-1}(H) = \frac{n}{2} - a(H) \Rightarrow \mu_{\frac{n}{2}}(G_n) \leq \frac{n}{2} + a(H),$$

which gives the result in (12) for even n . Otherwise, n is odd. First, we assume that $\mu_1(\overline{H}) \geq \left\lfloor \frac{n}{2} \right\rfloor$, that is, $a(H) \leq 1$. From this and (13), we have

$$\mu_{\frac{n-1}{2}}(\overline{G}_n) \geq \frac{n-1}{2} \Rightarrow \mu_{\frac{n+1}{2}}(G_n) \leq \frac{n+1}{2}.$$

Next, we assume that $\mu_1(\overline{H}) < \left\lfloor \frac{n}{2} \right\rfloor$, that is, $a(H) > 1$. From this and (13), we have

$$\mu_{\frac{n-1}{2}}(\overline{G}_n) \geq \mu_1(\overline{H}) \Rightarrow \mu_{\frac{n+1}{2}}(G_n) \leq \frac{n-1}{2} + a(H).$$

This completes the proof. \square

By similar arguments as those in Theorem 3.18, one can easily prove the following theorem on e -decomposable Riordan graphs.

Theorem 3.19. *Let $G_n = G_n(g, f)$ be an e -decomposable Riordan graph and let $H \cong \langle V_e \rangle$. Then*

$$\mu_{\lfloor \frac{n}{2} \rfloor+1}(G_n) \leq \left\lfloor \frac{n}{2} \right\rfloor \quad \text{and} \quad \mu_{\lfloor \frac{n}{2} \rfloor}(G_n) \leq \left\lfloor \frac{n}{2} \right\rfloor + a(H).$$

4 Vertex degrees in Riordan graphs

From now on, we assume that $p = \lfloor \log_2(n-1) \rfloor$ and $\{g\}_{-1}(1) = 0$. In the following, we obtain the degree of the vertex $2^p + 1$ in an io-decomposable Riordan graph $G_n(g, zg)$.

Lemma 4.1. *Let $G_n = G_n(g, zg)$ be an io-decomposable Riordan graph. Then*

$$d_{G_n}(2^p + 1) = 2^p + \{g\}_{n-2^p-2}(1). \quad (14)$$

In particular, the vertex $2^p + 1$ is universal in G_n if and only if $n = 2^p + 1$ or $n \neq 2^p + 1$ and $[z^{2^s}]g \equiv 1$ for $0 \leq s \leq \lfloor \frac{n-2^p}{2} \rfloor - 1$.

Proof: Suppose that $H = \langle \{2^p + 1, \dots, n\} \rangle$. Then

$$\begin{aligned} d_{G_n}(2^p + 1) &= d_{G_{2^p+1}}(2^p + 1) + d_H(2^p + 1) \\ &= d_{G_{2^p+1}}(2^p + 1) + d_{G_{n-2^p}}(1) \quad (\text{by Theorem 3.6(i) in [3]}) \\ &= 2^p + \{g\}_{n-2^p-2}(1) \quad (\text{by Theorems 4.13 and 3.1 in [3]}). \end{aligned}$$

This completes the proof for (14). The second part of the proof follows from (14) and Lemma 4.4 in [3]. \square

For any io-decomposable Riordan graph $G_n = G_n(g, zg)$, we have $[z^0]g = [z^1]g \equiv 1$. This observation, along with Lemma 14, gives the following result.

Corollary 4.2. *Let $G_n = G_n(g, zg)$ be an io-decomposable Riordan graph. Then the vertex $2^p + 1$ is universal in G_n for $2^p + 1 \leq n \leq 2^p + 3$.*

In the following two lemmas, some comparisons between the degree of the vertex $2^p + 1$ and degrees of other vertices in an io-decomposable Riordan graph are given.

Lemma 4.3. *Let $G_n = G_n(g, zg)$ be an io-decomposable Riordan graph. Then*

$$d_{G_n}(2^p + 1) \geq d_{G_n}(1). \quad (15)$$

Moreover, for $n \leq 1 + 2^{p-1} + 2^p$, we have

$$d_{G_n}(2^p + 1) \geq d_{G_n}(2^{p-1} + 1). \quad (16)$$

Proof: From Lemma 4.1, we have

$$d_{G_n}(2^p + 1) = 2^p + \{g\}_{n-2^p-2}(1) \geq \sum_{i=n-2^p-1}^{n-2} [z^i]g + \{g\}_{n-2^p-2}(1) = \{g\}_{n-2}(1) = d_{G_n}(1).$$

This completes the proof of (15).

Now, suppose that $n \leq 1 + 2^{p-1} + 2^p$. Then

$$\{g\}_{2^{p-1}-1}(1) - \{g\}_{n-2^p-2}(1) = \sum_{i=n-2^p-1}^{2^{p-1}-1} [z^i]g \leq 2^p + 2^{p-1} + 1 - n. \quad (17)$$

We define the subgraphs H_1 and H_2 of G_n as follows:

$$H_1 \cong \begin{cases} \langle \{2^{p-1} + 1, 2^p + 2, \dots, n\} \rangle & \text{if } n > 2^p + 1 \\ \langle \{2^{p-1} + 1\} \rangle & \text{if } n = 2^p + 1 \end{cases}$$

and $H_2 \cong \langle \{2^{p-1} + 1, \dots, 2^p + 1\} \rangle$. By Lemma 4.1, we have

$$\begin{aligned}
d_{G_n}(2^p + 1) &= 2^p + \{g\}_{n-2^{p-2}}(1) \\
&\geq n - 2^{p-1} - 1 + \{g\}_{2^{p-1}-1}(1) \quad (\text{by (17)}) \\
&= 2^{p-1} + n - 2^p - 1 + \{g\}_{2^{p-1}-1}(1) \\
&\geq 2^{p-1} + \{g\}_{2^{p-1}-1}(1) + d_{H_1}(2^{p-1} + 1) \\
&= d_{G_{2^{p-1}+1}}(2^{p-1} + 1) + d_{G_{2^{p-1}+1}}(1) + d_{H_1}(2^{p-1} + 1) \\
&= d_{G_{2^{p-1}+1}}(2^{p-1} + 1) + d_{H_2}(2^{p-1} + 1) + d_{H_1}(2^{p-1} + 1) \\
&= d_{G_n}(2^{p-1} + 1),
\end{aligned}$$

where the next to last equality is given by Theorem 3.6 in [3], and the last inequality is obtained using the fact that $d_{H_1}(2^{p-1} + 1) \leq n - 2^p - 1$. This completes the proof of (16). \square

Lemma 4.4. *Let $G_n = G_n(g, zg)$ be an io-decomposable Riordan graph. Then, for $i \in V_e$,*

$$d_{G_n}(2^p + 1) \geq d_{G_n}(i).$$

Proof: We have $n \leq 2^{p+1}$, that is, $\lceil \frac{n}{2} \rceil \leq 2^p$. From this and (14), for $i \in V_e$, we obtain

$$d_{G_n}(2^p + 1) \geq 2^p \geq \lceil \frac{n}{2} \rceil \geq d_{G_n}(i).$$

\square

Corollary 4.5. *Let $G_n = G_n(g, zg)$ be an io-decomposable Riordan graph. Then*

$$2^p + \lceil \log_2(n - 2^p) \rceil \leq d_{G_n}(2^p + 1) \leq n - 1.$$

Proof: By Lemma 4.1, we have

$$d_{G_n}(2^p + 1) = 2^p + \{g\}_{n-2^{p-2}}(1). \quad (18)$$

Since G_n is an io-decomposable Riordan graph, by Lemma 2.11, we have $g_{2^p-1} \equiv 1$ for $p \geq 0$. From this, (18) and Lemma 2.10, we obtain

$$d_{G_n}(2^p + 1) = 2^p + \{g\}_{n-2^{p-2}}(1) \geq 2^p + \lceil \log_2(n - 2^p) \rceil.$$

This completes our proof. \square

Lemma 4.6 ([3]). *Let $G_n(g, zg)$ be an io-decomposable Riordan graph. Then it is $(\lceil \log_2 n \rceil + 1)$ -partite with partitions $V_1, V_2, \dots, V_{\lceil \log_2 n \rceil + 1}$ such that*

$$\begin{aligned}
V_j &= \left\{ 2^{j-1} + 1 + i2^j \mid 0 \leq i \leq \left\lfloor \frac{n-1-2^{j-1}}{2^j} \right\rfloor \right\} \quad \text{if } 1 \leq j \leq \lceil \log_2 n \rceil, \\
V_{\lceil \log_2 n \rceil + 1} &= \{1\}.
\end{aligned}$$

It follows from Lemma 4.3 that if the vertex 1 is universal in an io-decomposable Riordan graph $G_n(g, zg)$, then the vertex $2^p + 1$ is also universal in that graph. In what follows, we show that if the vertex $2^{p-1} + 1$ is universal, then the vertex $2^p + 1$ is also universal.

Theorem 4.7. *Let $G_n = G_n(g, zg)$ be an io-decomposable Riordan graph. If $2^{p-1} + 1$ is universal then the vertex $2^p + 1$ is also universal.*

Proof: By Lemma 4.3, it suffices to prove that if $2^{p-1} + 1$ is a universal vertex in G_n , then $n \leq 1 + 2^{p-1} + 2^p$. By Lemma 4.6, we obtain that the vertex $2^{\lceil \log_2 n \rceil - 2} + 1 \in V_{\lceil \log_2 n \rceil - 1}$.

We claim that $p = \lceil \log_2 n \rceil - 1$. Indeed, by definition of p , we have $2^p + 1 \leq n \leq 2^{p+1}$. We may assume that $n = 2^p + \gamma$, where $1 \leq \gamma \leq 2^p$. So,

$$p = \lceil \log_2(2^p + 1) \rceil - 1 \leq \lceil \log_2 n \rceil - 1 = \lceil \log_2(2^p + \gamma) \rceil - 1 \leq \lceil \log_2 2^{p+1} \rceil - 1 = p,$$

which completes the proof of our claim.

Thus, $2^{p-1} + 1 \in V_p$. Again, by Lemma 4.6, we have $|V_p| = \left\lfloor \frac{n-1-2^{p-1}}{2^p} \right\rfloor + 1$. Since the vertex $2^{p-1} + 1$ is universal, we must have $|V_p| = 1$ and, consequently, $\left\lfloor \frac{n-1-2^{p-1}}{2^p} \right\rfloor = 0$, so $\frac{n-1-2^{p-1}}{2^p} < 1$ and $n < 1 + 2^{p-1} + 2^p$. This completes the proof. \square

Based on the facts above, we state the following conjecture.

Conjecture 4.8. *The vertex $2^p + 1$ has the maximum degree in any io-decomposable Riordan graph $G_n = G_n(g, zg)$.*

The following result is obtained from the facts above.

Corollary 4.9. *Let $G_n = G_n(g, zg)$ be an io-decomposable Riordan graph. Then G_n has at most three universal vertices which must come from the set $\{1, 2^{p-1} + 1, 2^p + 1\}$. In particular, if $n \geq 1 + 2^{p-1} + 2^p$, then G_n has at most two universal vertices that must come from the set $\{1, 2^p + 1\}$.*

The following result on Pascal graphs is obtained in [4].

Lemma 4.10 ([4]). *For a Pascal graph PG_n of order n , we have*

- (i) *If $n = 2^p + 1$, then the universal vertices are $1, \frac{n+1}{2}, n$.*
- (ii) *If $n \neq 2^p + 1$, then the universal vertices are $1, 2^p + 1$.*

By Corollary 4.9, for any io-decomposable Riordan graph of the Bell type with three universal vertices, the vertex 1 must be a universal vertex. The latter implies that $g(z) = \frac{1}{1-z}$ and $G_n \cong PG_n$, and thus, by Lemma 4.10, we have the following result.

Theorem 4.11. *Let $G_n = G_n(g, zg)$ be an io-decomposable Riordan graph with three universal vertices. Then $G_n \cong PG_n$ and $n = 2^p + 1$.*

Corollary 4.12. *Let $G_n = G_n(g, zg)$ be an io-decomposable Riordan graph with exactly one universal vertex. Then the universal vertex must be $2^p + 1$.*

Corollary 4.13. *Let $G_n = G_n(g, zg)$ be an io-decomposable Riordan graph. Then at most three Laplacian eigenvalues of G_n are equal to n . In particular, if $n \geq 1 + 2^{p-1} + 2^p$, then at most two Laplacian eigenvalues of G_n are equal to n .*

Theorems 4.14 and 4.15 below give bounds for $\mu_1(G_n)$ and $\mu_{n-1}(G_n)$, respectively, in terms of the generating function $g(z)$ for any io-decomposable Riordan graph of the Bell type.

Theorem 4.14. *Let $G_n = G_n(g, zg)$ be an io-decomposable Riordan graph. Then*

$$\mu_1(G_n) \geq 2^p + \{g\}_{n-2^p-2}(1) + 1. \quad (19)$$

The equality holds if and only if $n = 2^p + 1$, or $n \neq 2^p + 1$ and $[z^{2^s}]g \equiv 1$ for $0 \leq s \leq \lfloor \frac{n-2^p}{2} \rfloor - 1$.

Proof: By Lemmas 4.1 and 2.22, we obtain

$$\mu_1(G_n) \geq \Delta(G_n) + 1 \geq d(2^p + 1) + 1 = 2^p + \{g\}_{n-2^p-2}(1) + 1. \quad (20)$$

This completes the proof of (19). To prove the rest, we first assume that $n = 2^p + 1$. In this case, $\Delta(G_n) = n - 1$ and we obtain $\mu_1(G_n) \geq n$. From this and Lemma 2.20, we obtain $\mu_1(G_n) = n$. Otherwise, $n \neq 2^p + 1$ and $[z^{2^s}]g \equiv 1$ for $0 \leq s \leq \lfloor \frac{n-2^p}{2} \rfloor - 1$. From this, we obtain $\{g\}_{n-2^p-2}(1) = n - 2^p - 1$, and thus $2^p + \{g\}_{n-2^p-2}(1) + 1 = n$. This gives $\mu_1(G_n) \geq n$ and, again, by Lemma 2.20, we obtain $\mu_1(G_n) = n = 2^p + \{g\}_{n-2^p-2}(1) + 1$.

Conversely, we assume that $\mu_1(G_n) = 2^p + \{g\}_{n-2^p-2}(1) + 1$. By (20),

$$\mu_1(G_n) = \Delta(G_n) + 1 = d(2^p + 1) + 1.$$

But then, by Lemma 2.22, $d(2^p + 1) = \Delta(G_n) = n - 1$, and thus, by Lemma 4.1, we obtain $n = 2^p + 1$, or $n \neq 2^p + 1$ and $[z^{2^s}]g \equiv 1$ for $0 \leq s \leq \lfloor \frac{n-2^p}{2} \rfloor - 1$. We are done. \square

Theorem 4.15. *Let $G_n = G_n(g, zg)$ be an io-decomposable Riordan graph. Then*

$$\mu_{n-1}(G_n) \leq 1 + \sum_{i=1}^{\lfloor \frac{n+1}{2} \rfloor} g_{2i-1}.$$

Moreover, for $n \in \{2^p + 1, 2^p + 2\}$, we have $\mu_{n-1}(G_n) \geq 1$.

Proof: By Lemma 3.14, we obtain

$$d_{G_n}(2) = 1 + d_{G_{\lfloor \frac{n}{2} \rfloor}}(1) = 1 + \sum_{i=1}^{\lfloor \frac{n+1}{2} \rfloor} g_{2i-1}.$$

Since $G_n \not\cong K_n$, by Lemma 2.21, we obtain

$$\mu_{n-1}(G_n) \leq \delta(G_n) \leq d_{G_n}(2) = 1 + \sum_{i=1}^{\lfloor \frac{n+1}{2} \rfloor} g_{2i-1},$$

as desired. Now, for $n \in \{2^p + 1, 2^p + 2\}$, by Corollary 4.2, we have that G_n has at least one universal vertex. This means that the complement graph $\overline{G_n}$ of the io-decomposable Riordan graph G_n has at least one isolated vertex. Thus, $\mu_1(\overline{G_n}) \leq n - 1$ and $\mu_{n-1}(G) \geq 1$. \square

We end this section by using generating functions to study the *Rayleigh-Ritz quotient* for Riordan graphs. It is well known that for a graph G , the Rayleigh-Ritz quotient under the adjacency matrix $\mathcal{A}(G)$ provides a lower bound on the spectral radius $\lambda_1(G)$ of G as follows.

Lemma 4.16 ([2]). *Let G be a graph with n vertices and with the adjacency matrix $\mathcal{A}(G)$. For any nonzero vector $X = [x_1, x_2, \dots, x_n]^T \in \mathbb{R}^n$,*

$$\lambda_1(G) \geq \frac{X^t \mathcal{A}(G) X}{X^t X}.$$

The equality holds if and only if X is the eigenvector of $\mathcal{A}(G)$ with the spectral radius $\lambda_1(G)$.

Let $k = k(z) := k_0 + k_1 z + \dots + k_n z^n$ and $\ell = \ell(z) := \ell_0 + \ell_1 z + \dots + \ell_n z^n$ be two polynomials of degree n in $R[[z]]$. Similarly to the inner product between two vectors, we define

$$k \odot \ell = k(z) \odot \ell(z) := k_0 \ell_0 + k_1 \ell_1 + \dots + k_n \ell_n.$$

The following result gives a lower bound for the spectral radius of a Riordan graph of the Appell type.

Theorem 4.17. *Let $G_n = G_n(g, z)$ be a Riordan graph of the Appell type with n vertices. Then for any polynomial h of degree $n - 1$, we have*

$$\lambda_1(G_n) \geq \frac{2(h \odot k)}{h \odot h}, \quad (21)$$

where $k = \sum_{i=0}^{n-2} ([z^i]gh)z^i$. The equality in (21) holds if and only if h is the generating function of the eigenvector of G_n with eigenvalue $\lambda_1(G_n)$.

Proof: Let \mathcal{A}_n be the adjacency matrix of a Riordan graph $G_n(g, f)$. By Lemma 4.16 and (1), for any nonzero vector $X \in R^n$, we have

$$\begin{aligned} \lambda_1(G_n(g, f)) &\geq \frac{X^T \mathcal{A}_n X}{X^T X} = \frac{X^T ((zg, f)_n + (zg, f)_n^T) X}{X^T X} \\ &= \frac{X^T (zg, f)_n X + X^T (zg, f)_n^T X}{X^T X}. \end{aligned}$$

Since $(X^T (zg, f)_n X)^T = X^T (zg, f)_n^T X$ and $X^T (zg, f)_n X \in R$, we obtain

$$X^T (zg, f)_n X = X^T (zg, f)_n^T X \text{ and } \lambda_1(G_n(g, f)) \geq 2 \frac{X^T (zg, f)_n X}{X^T X}.$$

Now, we let $Y := (zg, f)_n X$ and $\ell(z)$ and $h(z)$ be, respectively, the generating functions of Y and X . Then, for $f = z$, we have

$$\lambda_1(G_n) \geq 2 \frac{X^T Y}{X^T X} = 2 \frac{h \odot \ell}{h \odot h} = 2 \frac{h \odot (zgh(f))}{h \odot h} = 2 \frac{h \odot (zgh)}{h \odot h},$$

where we have applied the FTRM (Lemma 2.1) and the fact that $zgf^j = z^{j+1}g \in Z_2[[z]]$ for any $0 \leq j \leq n - 2$. This completes the proof of (21). The second part of the proof follows directly from the equality condition in Lemma 4.16. \square

Next, for the sake of example, we demonstrate that Theorem 4.17 works for complete graphs.

Example 4.18. *It is well known that for a complete graph K_n , $\lambda_1(K_n) = n - 1$ with eigenvector $X^T = \underbrace{[1, \dots, 1]}_n$. The generating function of X is $h(z) = \sum_{i=0}^{n-1} z^i$. Moreover, by Theorem 4.17,*

for $K_n \cong G_n\left(\frac{1}{1-z}, z\right)$, we have

$$h \odot h = \sum_{i=0}^{n-1} 1 = n, \quad k(z) = \sum_{i=1}^{n-1} iz^{i-1}, \quad h \odot k = \sum_{i=1}^{n-1} i = \frac{n(n-1)}{2},$$

and hence, by Theorem 4.17, indeed $\lambda_1(K_n) = n - 1$.

5 Eigenvalues of Pascal graphs and Catalan graphs

The following conjecture shows the significance of Pascal graphs PG_n and Catalan graphs CG_n , and we devote this section to studying eigenvalues of these graphs.

Conjecture 5.1 ([3]). *Let G_n be an io-decomposable Riordan graph of the Bell type. Then,*

$$2 = \text{diam}(PG_n) \leq \text{diam}(G_n) \leq \text{diam}(CG_n)$$

for $n \geq 4$. Moreover, PG_n is the only graph in the class of io-decomposable graphs of the Bell type whose diameter is 2 for all $n \geq 4$.

In this section, we first present some results related to eigenvalues -1 and 0 of Pascal and Catalan graphs, and next, we give some results about integral Laplacian eigenvalues of these graphs.

The following result can be directly proved by definition of an eigenvalue and a corresponding eigenvector of a graph.

Lemma 5.2. *Let G be a graph of order n . If $uv \in E(G)$ with $N_G(u) \setminus \{v\} = N_G(v) \setminus \{u\}$, then -1 is an eigenvalue of G with eigenvector*

$$X = [0, \dots, 0, 1, 0, \dots, 0, -1, 0, \dots, 0]^T,$$

where the entries 1 and -1 correspond to the vertices u and v , respectively.

Recall that the Catalan graph CG_n is defined by $G_n \left(\frac{1-\sqrt{1-4z}}{2z}, \frac{1-\sqrt{1-4z}}{2} \right)$.

Theorem 5.3. *Let $n \geq 2$. Then -1 is an eigenvalue of a Catalan graph CG_n with eigenvector $X = [1, -1, 0, \dots, 0]^T$.*

Proof: We know that $CG_n = G_n(g, zg)$, where $g = \frac{1-\sqrt{1-4z}}{2z}$. Suppose that $C_1(z)$ and $C_2(z)$ are the generating functions for the first and the second columns of the binary Riordan matrix $\mathcal{B}(zg, zg)_n$. We have that $C_1(z) = zg = \frac{1-\sqrt{1-4z}}{2}$ and $C_2(z) = zgf = z^2g^2 = \frac{2-2\sqrt{1-4z}-4z}{4} = \frac{1-\sqrt{1-4z}}{2} - z$. From this, using the fact that $12 \in E(CG_n)$, we obtain $N_{CG_n}(1) \setminus \{2\} = N_{CG_n}(2) \setminus \{1\}$. Now, by Lemma 5.2, the desired result is obtained. \square

The following result is an immediate corollary of Lemmas 5.2 and 4.10.

Theorem 5.4. *For a Pascal graph PG_n , we have*

- (i) *If $n = 2^p + 1$, then -1 is an eigenvalue of PG_n with multiplicity at least 2 and with eigenvectors $X = [1, \underbrace{0, \dots, 0}_{\frac{n-1}{2}}, -1, 0, \dots, 0]^T$ and $Y = [1, 0, \dots, 0, -1]^T$.*
- (ii) *If $n \neq 2^p + 1$, then -1 is an eigenvalue of PG_n with eigenvector $X = [1, \underbrace{0, \dots, 0}_{2^p}, -1, 0, \dots, 0]^T$.*

Lemma 5.5 ([8]). *Let G be a graph of order n . If $uv \notin E(G)$ and $N_G(u) = N_G(v)$ then 0 is an eigenvalue of G with eigenvector*

$$X = [0, \dots, 0, 1, 0, \dots, 0, -1, 0, \dots, 0]^T,$$

where the entries 1 and -1 correspond to the vertices u and v , respectively.

Theorem 5.6. *Let $2^p \leq n \leq 2^p + 2$. Then 0 is an eigenvalue of a Pascal graph PG_n with eigenvector X , where the entries of X in positions 2 and 2^p are 1 and -1 , respectively, and the other entries are 0 s.*

Proof: From [4], we have $N_{PG_n}(2) = V_o$ and $N_{PG_n}(2^p) = \{j \in V_o \mid 1 \leq j \leq 2^p + 1\}$. Since $2^p \leq n \leq 2^p + 2$, we have $N_{PG_n}(2) = N_{PG_n}(2^p)$. Finally, by the fact that $2(2^p) \notin E(PG_n)$ and Lemma 5.5, we obtain the desired. \square

Now, we are ready to present some results on integral Laplacian eigenvalues of Pascal and Catalan graphs. The following result is obtained by Lemma 3.1 in [6].

Lemma 5.7. *Let G be a graph of order n . If $uv \notin E(G)$ and $N_G(u) = N_G(v)$, then $d_G(u) = |N_G(u)|$ is a Laplacian eigenvalue of G with eigenvector*

$$X = [0, \dots, 0, 1, 0, \dots, 0, -1, 0, \dots, 0]^T,$$

where the entries 1 and -1 correspond to the vertices u and v , respectively.

Theorem 5.8. *Let $2^p \leq n \leq 2^p + 2$. Then $\lceil \frac{n}{2} \rceil$ is a Laplacian eigenvalue of a Pascal graph PG_n with eigenvector X , where the entries in positions 2 and 2^p are 1 and -1 , respectively, and the other entries are 0s.*

Proof: By the proof of Theorem 5.6, we have $N_{PG_n}(2) = N_{PG_n}(2^p)$. Since $2(2^p) \notin E(PG_n)$, the required result is obtained by Lemma 5.7. \square

The following result is obtained by Theorem 3.3 in [7].

Lemma 5.9. *Let G be a graph of order n . If $uv \in E(G)$ with $N_G(u) \setminus \{v\} = N_G(v) \setminus \{u\}$, then $d_G(u) + 1$ is a Laplacian eigenvalue of G with eigenvector*

$$X = [0, \dots, 0, 1, 0, \dots, 0, -1, 0, \dots, 0]^T,$$

where the entries 1 and -1 correspond to the vertices u and v , respectively.

The following result is related to Laplacian eigenvalues of Catalan graphs.

Theorem 5.10. *Let $n \geq 2$. Then $\lceil \log_2 n \rceil + 1$ is a Laplacian eigenvalue of a Catalan graph CG_n with eigenvector $X = [1, -1, 0, \dots, 0]^T$.*

Proof: By the arguments used in the proof of Theorem 5.3, we have

$$N_{CG_n}(1) \setminus \{2\} = N_{CG_n}(2) \setminus \{1\}.$$

Now, by Lemmas 2.23 and 5.9, the desired result is obtained. \square

The following result follows directly from Lemma 4.10 and Theorem 3.3 in [7].

Theorem 5.11. *For a Pascal graph PG_n , we have*

(i) *If $n = 2^p + 1$, then n is a Laplacian eigenvalue of PG_n with multiplicity at least 2 and with eigenvectors $X = [1, \underbrace{0, \dots, 0}_{\frac{n-1}{2}}, -1, 0, \dots, 0]^T$ and $Y = [1, 0, \dots, 0, -1]^T$.*

(ii) *If $n \neq 2^p + 1$, then n is a Laplacian eigenvalue of PG_n with eigenvector*

$$X = [1, \underbrace{0, \dots, 0}_{2^p}, -1, 0, \dots, 0]^T.$$

We note that Theorem 4.3 in [4] gives a better result than Theorem 5.11 about the number of Laplacian eigenvalues equal to n , but without any information about the eigenvectors.

6 The inertias of Riordan graphs

In this section, we present some results on the positive inertia and the negative inertia of Riordan graphs. We begin with two lemmas.

Lemma 6.1. *Let $G_n = G_n(g, f)$ be a Riordan graph with n vertices. Then, the following are equivalent:*

(i) $[z^{2k}](gf) \equiv 0$ for $1 \leq k \leq \lfloor \frac{n-2}{2} \rfloor$.

(ii) $2j \notin E(G_n)$ for $j \in \{4, 6, \dots, 2 \lfloor \frac{n}{2} \rfloor\}$.

(iii) G_n is o -decomposable.

Proof: The statements (i) and (ii) are clearly equivalent. We have (i) holds if and only if

$$\left(\frac{gf}{z}\right)'(\sqrt{z}) = \sum_{k=1}^{\lfloor \frac{n-2}{2} \rfloor} \left(\sum_{i=1}^{2k} g_{2k-i} f_i \right) z^{2k} = \sum_{k=1}^{\lfloor \frac{n-2}{2} \rfloor} ([z^{2k}](gf)) z^{2k} \equiv 0.$$

By Theorem 2.4, this is true if and only if $Y = O_{\lfloor \frac{n}{2} \rfloor}$, i.e. $\langle V_e \rangle \cong N_{\lfloor \frac{n}{2} \rfloor}$, and G_n is e-decomposable by definition. \square

Lemma 6.2. *Let $G_n = G_n(g, f)$ be a Riordan graph with n vertices. Then, the following are equivalent:*

- (i) $[z^{2i-1}]g \equiv 0$ for $1 \leq i \leq \lfloor \frac{n-1}{2} \rfloor$.
- (ii) $1j \notin E(G_n)$ for $j \in \{3, 5, \dots, 2\lfloor \frac{n}{2} \rfloor - 1\}$.
- (iii) G_n is e-decomposable.

Proof: The statements (i) and (ii) are clearly equivalent. Further, we have that (i) holds if and only if

$$g'(\sqrt{z}) = \sum_{i=1}^{\lfloor \frac{n-1}{2} \rfloor} g_{2i-1} z^{2i-1} \equiv 0.$$

By Theorem 2.4, this is true if and only if $X = O_{\lfloor \frac{n}{2} \rfloor}$, i.e. $\langle V_o \rangle \cong N_{\lfloor \frac{n}{2} \rfloor}$, and G_n is e-decomposable by definition. \square

The following result is obtained by Lemmas 6.2 and 6.1.

Corollary 6.3. *Let G_n be a Riordan graph with n vertices.*

- (i) *If $G_n = G_n(g, zg)$ is of the Bell type, then G_n is o-decomposable.*
- (ii) *If $G_n = G_n(f', f)$ is of the derivative type, then G_n is e-decomposable.*

Now, using the lemmas above, we obtain the following results on the positive and negative inertias of Riordan graphs.

Theorem 6.4. *Let G_n be an o-decomposable Riordan graph. Then,*

$$\max\{n^+, n^-\} \leq \left\lceil \frac{n}{2} \right\rceil.$$

Proof: By Lemma 6.1, the null graph $N_{\lfloor \frac{n}{2} \rfloor}$ is an induced subgraph of G_n . This fact, and the interlacing property in Lemma 2.14, gives

$$\lambda_i(G_n) \geq \lambda_i\left(N_{\lfloor \frac{n}{2} \rfloor}\right) \geq \lambda_{\lfloor \frac{n}{2} \rfloor + i}(G_n) \quad \text{for } 1 \leq i \leq \left\lfloor \frac{n}{2} \right\rfloor. \quad (22)$$

Putting $i = \lfloor \frac{n}{2} \rfloor$ in the first inequality of (22), we obtain

$$\lambda_{\lfloor \frac{n}{2} \rfloor}(G_n) \geq \lambda_{\lfloor \frac{n}{2} \rfloor}\left(N_{\lfloor \frac{n}{2} \rfloor}\right) = 0,$$

that is, $n^+ + \eta \geq \lfloor \frac{n}{2} \rfloor$, which implies that $n^- \leq \lceil \frac{n}{2} \rceil$. Now, putting $i = 1$ in the second inequality of (22), we obtain

$$\lambda_{\lfloor \frac{n}{2} \rfloor + 1}(G_n) \leq \lambda_1\left(N_{\lfloor \frac{n}{2} \rfloor}\right) = 0,$$

that is, $n^- + \eta \geq \lfloor \frac{n}{2} \rfloor$, which implies $n^+ \leq \lceil \frac{n}{2} \rceil$. We are done. \square

Corollary 6.5. *Let G_n be an o -decomposable Riordan graph. If n is even and $\det(G_n) \neq 0$, then $n^+ = n^- = \frac{n}{2}$.*

Corollary 6.6. *Let $G_n = G_n(g, zg)$ be a Riordan graph of the Bell type with n vertices. Then,*

$$\max\{n^+, n^-\} \leq \left\lceil \frac{n}{2} \right\rceil. \quad (23)$$

Moreover, if n is even and $\det(G_n) \neq 0$, then $n^+ = n^- = \frac{n}{2}$.

The following result is obtained using similar arguments as those in the proofs of Theorem 6.4 and Lemma 6.2.

Theorem 6.7. *Let G_n be an e -decomposable Riordan graph. Then,*

$$\max\{n^+, n^-\} \leq \left\lfloor \frac{n}{2} \right\rfloor.$$

Next, we state two corollaries of Theorem 6.7.

Corollary 6.8. *Let G_n be an e -decomposable Riordan graph. We have the following:*

- (i) *If n is odd, then $\det(G_n) = 0$.*
- (ii) *If $\det(G_n) \neq 0$, then n is even and $n^+ = n^- = \frac{n}{2}$.*

Corollary 6.9. *Let $G_n = G_n(f', f)$ be a Riordan graph of the derivative type. We have the following:*

- (i) $\max\{n^+, n^-\} \leq \lfloor \frac{n}{2} \rfloor$.
- (ii) *If n is odd, then $\det(G_n) = 0$.*
- (iii) *If $\det(G_n) \neq 0$, then n is even and $n^+ = n^- = \frac{n}{2}$.*

Example 6.10. *In what follows, we used Sage [16] to calculate n^+ and n^- .*

- (i) *The Pascal graph PG_{10} is a Riordan graph of the Bell type. We have $n^+(PG_{10}) + 1 = n^-(PG_{10}) = 5$. This is consistent with Corollary 6.6 and gives the equality in (23).*
- (ii) *The graph $G_{10}\left(\frac{1}{1+z^2}, \frac{z}{1+z}\right)$ is a Riordan graph of the derivative type. We have $n^+\left(G_{10}\left(\frac{1}{1+z^2}, \frac{z}{1+z}\right)\right) = n^-\left(G_{10}\left(\frac{1}{1+z^2}, \frac{z}{1+z}\right)\right) = 5$. This is consistent with Corollary 6.9 and gives the equality in (i).*
- (iii) $n^+\left(G_{16}\left(1+z^3, \frac{z}{1+z}\right)\right) = 6$, $n^-\left(G_{16}\left(1+z^3, \frac{z}{1+z}\right)\right) = 10$.
- (iv) $n^+(K_n) = 1$, $n^-(K_n) = n - 1$.

Clearly, $G_{16}\left(1+z^3, \frac{z}{1+z}\right)$ and K_n do not satisfy the conditions in Lemmas 6.2 and 6.1.

Using Sage [16], we observe that $n^-(PG_i) = \lceil \frac{i}{2} \rceil$ for $2 \leq i \leq 200$. Moreover, by Lemma 2.6, any Riordan graph $G_n = G_n(g, zg)$ with an even function g is bipartite and thus $n^+(G_n) = n^-(G_n)$. Based on these observation, we state the following problem.

Problem 6.11. *Is it true that $n^+ \leq n^-$ for any Riordan graph $G_n(g, zg)$?*

Let H_n be the bipartite graph obtained from a Riordan graph $G_n(g, f)$ by deleting all edges in $\langle V_o \rangle$ and $\langle V_e \rangle$. The following result is obtained directly from Lemma 2.17.

Theorem 6.12. *Let $G_n = G_n(g, f)$ be an o-decomposable, or an e-decomposable, Riordan graph. Then,*

$$\min(n^+(G_n), n^-(G_n)) \geq \frac{1}{2} \text{rank}(H_n) = n^+(H_n) = n^-(H_n).$$

In particular, if n is even and the corresponding bipartite graph is nonsingular, then $n^-(G_n) = n^+(G_n) = n/2$ and $\eta(G_n) = 0$.

For a graph G , we denote the number of eigenvalues of G located in the interval I by $m_G(I)$. The following results give bounds on $m_{\overline{G}_n}(-\infty, -1]$.

Theorem 6.13. *Let $G_n = G_n(g, f)$ be an o-decomposable Riordan graph and \overline{G}_n be its complement. Then,*

$$n^-(\overline{G}_n) \geq m_{\overline{G}_n}(-\infty, -1] \geq \left\lfloor \frac{n}{2} \right\rfloor - 1.$$

Proof: From the definition of an o-decomposable Riordan graph, we have $\overline{K}_{\lfloor n/2 \rfloor}$ is an induced subgraph of G_n and, consequently, the complete graph $K_{\lfloor n/2 \rfloor}$ is an induced subgraph of \overline{G}_n . Now, by the well known interlacing theorem, we obtain

$$-1 = \lambda_2(K_{\lfloor n/2 \rfloor}) \geq \lambda_{n - \lfloor n/2 \rfloor + 2}(\overline{G}_n) = \lambda_{\lfloor n/2 \rfloor + 2}(\overline{G}_n),$$

which gives the desired result. \square

By a similar argument, one can easily obtain the following result.

Theorem 6.14. *Let $G_n = G_n(g, f)$ be an e-decomposable Riordan graph and \overline{G}_n be its complement. Then,*

$$n^-(\overline{G}_n) \geq m_{\overline{G}_n}(-\infty, -1] \geq \left\lceil \frac{n}{2} \right\rceil - 1.$$

7 The nullity of Riordan graphs

In this section, we present some results on the nullity of Riordan graphs.

Theorem 7.1. *Let \mathcal{A}_n be the adjacency matrix of an o-decomposable Riordan graph $G_n(g, f)$ of the form $P \begin{bmatrix} X_{\lceil \frac{n}{2} \rceil} & B \\ B^T & O \end{bmatrix} P^T$. Then*

$$\eta(B) \leq \eta(\mathcal{A}_n) \leq \begin{cases} 2\eta(B) & \text{if } n \text{ is even} \\ 2\eta(B) + 1 & \text{if } n \text{ is odd.} \end{cases}$$

Proof: We have

$$\text{rank}(B) + \text{rank}(B^T) = \text{rank} \left(\begin{bmatrix} O_{\lceil \frac{n}{2} \rceil} & B \\ B^T & O \end{bmatrix} \right) \leq \text{rank}(\mathcal{A}_n) \leq \text{rank} \left(\begin{bmatrix} X_{\lceil \frac{n}{2} \rceil} \\ B^T \end{bmatrix} \right) + \text{rank}(B).$$

Since $\text{rank}(B) = \text{rank}(B^T)$, from the above, we obtain

$$2 \text{rank}(B) \leq \text{rank}(\mathcal{A}_n) \leq \left\lceil \frac{n}{2} \right\rceil + \text{rank}(B).$$

We can now apply the rank-nullity theorem to obtain

$$2 \left(\left\lfloor \frac{n}{2} \right\rfloor - \eta(B) \right) \leq n - \eta(\mathcal{A}_n) \leq \left\lceil \frac{n}{2} \right\rceil + \left(\left\lfloor \frac{n}{2} \right\rfloor - \eta(B) \right) \Rightarrow \eta(B) \leq \eta(\mathcal{A}_n) \leq 2\eta(B) + n - 2 \left\lfloor \frac{n}{2} \right\rfloor.$$

This completes our proof. \square

Corollary 7.2. *Let $G_n(g, f)$ be an o -decomposable Riordan graph of even order n with the adjacency matrix \mathcal{A}_n of the form in Theorem 7.1. Then $\det B \neq 0$ if and only if $\det \mathcal{A}_n \neq 0$.*

Corollary 7.3. *Let $G_n(g, f)$ be an o -decomposable Riordan graph of odd order n with the adjacency matrix \mathcal{A}_n of the form in Theorem 7.1. If $\eta(B) = 0$ then $\eta(\mathcal{A}_n) = 0$ or 1.*

When the graph $G_n(g, zg)$ is proper (recall Definition 2.3), we have the following result on the matrix $P^T \mathcal{A}_n P$, where $\mathcal{A}_n = \mathcal{A}_n(g, zg)$. This result is obtained by Lemma 2.5.

Corollary 7.4. *Let $[z^0]g \equiv 1$ and $\mathcal{A}_n = \mathcal{A}_n(g, zg)$. Then in the matrix $P^T \mathcal{A}_n P$ in Lemma 3.14, we have the following.*

(i) *If n is odd, then*

$$b_{i,i} = 1 \quad \text{for } 1 \leq i \leq \frac{n-1}{2}, \quad \text{and} \quad r_{j,(j-1)} = 1 \quad \text{for } 2 \leq j \leq \frac{n+1}{2}.$$

(ii) *If n is even, then*

$$b_{i,i} = 1 \quad \text{for } 1 \leq i \leq \frac{n}{2}, \quad \text{and} \quad r_{j,(j-1)} = 1 \quad \text{for } 2 \leq j \leq \frac{n}{2}.$$

Theorem 7.5. *Let $[z^0]g \equiv 1$ and let*

$$\mathcal{A}_n := P \begin{bmatrix} \mathcal{A}_{\lceil \frac{n}{2} \rceil} & B \\ B^T & O \end{bmatrix} P^T$$

be the adjacency matrix of an io -decomposable Riordan graph $G_n(g, zg)$ of the Bell type. For any $n \geq 2$, we have

$$\text{rank} \left(\begin{bmatrix} \mathcal{A}_{\lceil \frac{n}{2} \rceil} \\ B^T \end{bmatrix} \right) = \left\lceil \frac{n}{2} \right\rceil.$$

Proof: We consider the cases of even and odd n separately.

(i) Let $n = 2k$ and

$$\begin{bmatrix} \mathcal{A}_k \\ B^T \end{bmatrix} = \begin{bmatrix} \mathbf{r}_1 & \cdots & \mathbf{r}_k & \mathbf{b}_1 & \cdots & \mathbf{b}_k \end{bmatrix}^T$$

where \mathbf{r}_i is the i th row of \mathcal{A}_k and \mathbf{b}_i is the i th row of B^T . We consider the $k \times k$ matrix

$$X_k = \begin{bmatrix} \mathbf{b}_1 - \mathbf{r}_1 & \cdots & \mathbf{b}_{k-1} - \mathbf{r}_{k-1} & \mathbf{b}_k \end{bmatrix}^T.$$

It follows from Lemma 3.14 and Remark 7.4 that X_k is a unit lower triangular matrix. Thus,

$$\text{rank} \left(\begin{bmatrix} \mathcal{A}_{\lceil \frac{n}{2} \rceil} \\ B^T \end{bmatrix} \right) = \text{rank}(X_k) = k = \left\lceil \frac{n}{2} \right\rceil.$$

(ii) Let $n = 2k + 1$. Note that $\mathcal{A}_{\lceil \frac{n}{2} \rceil} = \mathcal{A}_{k+1}$ and B^T is the $k \times (k+1)$ matrix. Using the same notation as above, we have

$$\begin{bmatrix} \mathcal{A}_{k+1} \\ B^T \end{bmatrix} = \begin{bmatrix} \mathbf{r}_1 & \cdots & \mathbf{r}_{k+1} & \mathbf{b}_1 & \cdots & \mathbf{b}_k \end{bmatrix}^T.$$

We consider the $(k+1) \times (k+1)$ matrix

$$Y_{k+1} = \begin{bmatrix} \mathbf{b}_1 - \mathbf{r}_1 & \cdots & \mathbf{b}_k - \mathbf{r}_k & \mathbf{r}_k \end{bmatrix}^T.$$

It follows from Lemma 3.14 and Remark 7.4 that Y_{k+1} is also a unit lower triangular matrix. Thus,

$$\text{rank} \left(\begin{bmatrix} \mathcal{A}_{\lceil \frac{n}{2} \rceil} \\ B^T \end{bmatrix} \right) = \text{rank}(Y_{k+1}) = k + 1 = \lceil \frac{n}{2} \rceil,$$

which completes our proof. \square

Let $X = [x_1, x_2, \dots, x_n]^T$ be an eigenvector of the matrix \mathcal{A}_n with eigenvalue 0. We define X_o and X_e to be the vectors of all eigencomponents x_i , respectively, with odd and even indices i , that is,

$$X_o = [x_1, x_3, \dots, x_{2\lceil \frac{n}{2} \rceil - 1}]^T \quad \text{and} \quad X_e = [x_2, x_4, \dots, x_{2\lfloor \frac{n}{2} \rfloor}]^T.$$

Lemma 7.6. *Let \mathcal{A}_n be the adjacency matrix of a singular io-decomposable Riordan graph $G_n(g, zg)$ of the form in Theorem 7.5. Then*

$$X_o = O_{\lceil \frac{n}{2} \rceil} \quad \text{if and only if} \quad BX_e = O_{\lceil \frac{n}{2} \rceil}.$$

Proof: By Theorem 2.5 and our assumptions, we have

$$A_n X = \begin{bmatrix} \mathcal{A}_{\lceil \frac{n}{2} \rceil} & B \\ B^T & O \end{bmatrix} \begin{bmatrix} X_o \\ X_e \end{bmatrix} = O_n, \quad (24)$$

$$\begin{aligned} \mathcal{A}_{\lceil \frac{n}{2} \rceil} X_o + BX_e &= O_{\lceil \frac{n}{2} \rceil}, \\ B^T X_o &= O_{\lfloor \frac{n}{2} \rfloor}. \end{aligned}$$

If $X_o = O_{\lceil \frac{n}{2} \rceil}$, obviously, we obtain $BX_e = O_{\lceil \frac{n}{2} \rceil}$. Now, we assume that $BX_e = O_{\lceil \frac{n}{2} \rceil}$. Then, $DX_o = O_n$, where

$$D = \begin{bmatrix} \mathcal{A}_{\lceil \frac{n}{2} \rceil} \\ B^T \end{bmatrix}.$$

Since, by Theorem 7.5, the matrix D is full column rank, we obtain $X_o = O_{\lceil \frac{n}{2} \rceil}$, as desired. \square

Using Sage [16], we come up with the following conjecture.

Conjecture 7.7. *Let $G_n(g, zg)$ be a singular io-decomposable Riordan graph. Then $X_o = O_{\lceil \frac{n}{2} \rceil}$.*

In what follows, we show that Conjecture 7.7 is equivalent to a condition on $\eta(G_n)$.

Theorem 7.8. *Let \mathcal{A}_n be the adjacency matrix of a singular io-decomposable Riordan graph $G_n(g, zg)$ of the form in Theorem 7.5. Then $X_o = O_{\lceil \frac{n}{2} \rceil}$ if and only if $\eta(G_n) = \eta(B)$.*

Proof: First, assume that $X_o = O_{\lceil \frac{n}{2} \rceil}$. Then, by Lemma 7.6, this is equivalent to $BX_e = O_{\lceil \frac{n}{2} \rceil}$. This leads to $\eta(G_n) = \eta(B)$.

Now, we assume that $\eta(G_n) = \eta(B)$, that is, that $\text{rank}(\mathcal{A}_n) = \lceil \frac{n}{2} \rceil + \text{rank}(B)$. Since X is an eigenvector of G_n with eigenvalue 0, we have

$$\begin{bmatrix} \mathcal{A}_{\lceil \frac{n}{2} \rceil} & B \\ B^T & O \end{bmatrix} \begin{bmatrix} X_o \\ X_e \end{bmatrix} = O_n.$$

Suppose that $\text{rank}(B) = r_b$. Using row operations on B , one can easily obtain $\begin{pmatrix} B_{r_b} \\ O \end{pmatrix}$, where B_{r_b} is an $r_b \times \lfloor \frac{n}{2} \rfloor$ matrix containing r_b linearly independent rows of B . Using the same row operations on $P^T \mathcal{A} P$, we have

$$\left[\begin{array}{c|c} A_1 & B_{r_b} \\ \hline A_2 & O \\ \hline B^T & O \end{array} \right] \begin{bmatrix} X_o \\ X_c \end{bmatrix} = O_n,$$

where A_2 is an $(\lfloor \frac{n}{2} \rfloor - r_b) \times \lfloor \frac{n}{2} \rfloor$ matrix. From this, we obtain $\begin{bmatrix} A_2 \\ B^T \end{bmatrix} X_o = O_k$, where $k = \lfloor \frac{n}{2} \rfloor + \lfloor \frac{n}{2} \rfloor - r_b$. To prove $X_o = O_{\lfloor \frac{n}{2} \rfloor}$, it suffices to show that

$$\text{rank} \left(\begin{bmatrix} A_2 \\ B^T \end{bmatrix} \right) = \lfloor \frac{n}{2} \rfloor. \quad (25)$$

We have

$$\lfloor \frac{n}{2} \rfloor + \text{rank}(B) = \text{rank} \left[\begin{array}{c|c} A_1 & B_{r_b} \\ \hline A_2 & O \\ \hline B^T & O \end{array} \right] \leq \text{rank}(B_r) + \text{rank} \left(\begin{bmatrix} A_2 \\ B^T \end{bmatrix} \right),$$

which gives the desired result in (25). Hence, the proof is done. \square

Applying Theorem 7.8, we can state a different version of Conjecture 7.7.

Conjecture 7.9. *Let \mathcal{A}_n be the adjacency matrix of a singular io-decomposable Riordan graph $G_n(g, zg)$ of the form in Theorem 7.5. Then, $\eta(G_n) = \eta(B)$.*

Lemma 7.10 ([1]). *Let A, B, C and D be, respectively, real $n \times n$, $n \times m$, $l \times n$ and $l \times m$ matrices, and let A be nonsingular. Then*

$$\text{rank} \begin{bmatrix} A & B \\ C & D \end{bmatrix} = n + \text{rank}(D - CA^{-1}B).$$

Theorem 7.11. *Let \mathcal{A}_n be the adjacency matrix of an io-decomposable Riordan graph $G_n(g, zg)$ of the form in Theorem 7.5. Then $\eta(G_n) = \eta(B^T M^{-1} B)$, where $M = \mathcal{A}_{\frac{n}{2}} - B^T$ for even n , and*

$$M = \begin{bmatrix} \mathbf{r}_1 - \mathbf{b}_1 & \cdots & \mathbf{r}_{k-1} - \mathbf{b}_{k-1} & \mathbf{r}_k - \mathbf{b}_{k-1} \end{bmatrix}^T$$

for odd n , $k = \frac{n+1}{2}$, where \mathbf{r}_i is the i th row of \mathcal{A}_k and \mathbf{b}_i is the i th row of B^T .

Proof: We know that the rank of any matrix is fixed under row and column operations. Thus, we have

$$\text{rank}(\mathcal{A}_n) = \text{rank} \begin{bmatrix} \mathcal{A}_{\lfloor \frac{n}{2} \rfloor} & B \\ B^T & O \end{bmatrix} = \text{rank} \begin{bmatrix} M_{\lfloor \frac{n}{2} \rfloor} & B \\ B^T & O \end{bmatrix}.$$

By Lemma 3.14, the square matrix M is a lower triangular matrix with all entries on the main diagonal equal to -1 . Therefore, M is invertible, and the required result is given by Lemma 7.10. \square

By Theorem 7.11, it is worth noticing that Conjecture 7.9 is true for any io-decomposable Riordan graph $G_n(g, zg)$ if and only if $\eta(B) = \eta(B^T M^{-1} B)$.

8 Determinants of Riordan graphs

In this section, we study determinants of o-decomposable and e-decomposable Riordan graphs, as well as determinants of Catalan graphs.

We begin with Schur's formula for determinant of a block matrix.

Lemma 8.1 ([1]). *Let A, B, C and D be real $n \times n$ matrices. Then*

$$\det \begin{bmatrix} A & B \\ C & D \end{bmatrix} = \begin{cases} \det(DA - CB), & \text{if } AB=BA; \\ \det(AD - BC), & \text{if } DC=CD. \end{cases}$$

Recall that H_n is the bipartite graph obtained from a Riordan graph $G_n(g, f)$ by deleting all edges in $\langle V_o \rangle$ and $\langle V_e \rangle$.

Theorem 8.2. *Let $G_n = G_n(g, f)$ be an o-decomposable, or e-decomposable, Riordan graph of even order n . Then,*

$$\det(G_n) = (\det(B))^2,$$

where B is given in (3).

Proof: First, suppose that \mathcal{A}_n is the adjacency matrix of an o-decomposable Riordan graph G_n . We have

$$\mathcal{A}_n := P \begin{bmatrix} X_{\lceil \frac{n}{2} \rceil} & B \\ B^T & O \end{bmatrix} P^T.$$

Since n is even and B is a square matrix, by Lemma 8.1, we obtain

$$\det(G_n) = \det(\mathcal{A}_n) = \det \begin{bmatrix} X_{\lceil \frac{n}{2} \rceil} & B \\ B^T & O \end{bmatrix} = \det(BB^T) = (\det(B))^2.$$

This completes the proof for o-decomposable Riordan graphs.

Next, suppose that \mathcal{A}_n is the adjacency matrix of an e-decomposable Riordan graph G_n . We have

$$\mathcal{A}_n := P \begin{bmatrix} O & B \\ B^T & Y_{\lfloor \frac{n}{2} \rfloor} \end{bmatrix} P^T.$$

Since n is even, B is a square matrix, and thus, by Lemma 8.1, we obtain

$$\det(G_n) = \det(\mathcal{A}_n) = \det \begin{bmatrix} O & B \\ B^T & Y_{\lfloor \frac{n}{2} \rfloor} \end{bmatrix} = \det(B^T B) = (\det(B))^2.$$

This completes the proof for e-decomposable Riordan graphs. □

The following results are obtained by Theorem 8.2.

Corollary 8.3. *Let $G_n = G_n(g, f)$ be an o-decomposable Riordan graph of even order n . Then $\det(B) = 0$ if and only if $\det(G_n) = 0$, that is, G_n is a singular graph if and only if H_n is such a graph. In particular, if $N_{H_n}(i) = N_{H_n}(j)$ for $i, j \in V_o$, $i \neq j$, then $\det(G_n) = 0$.*

Proof: Our first claim is a direct corollary of Theorem 8.2. Now, suppose that \mathbf{b}_i and \mathbf{b}_j are the i th and j th rows of the submatrix B . From our assumption, we obtain $\mathbf{b}_i = \mathbf{b}_j$, where $i, j \in V_o$ ($i \neq j$). This gives $\det(B) = 0$, and hence, by Theorem 8.2, the proof is complete. □

Corollary 8.4. *Let $G_n = G_n(g, f)$ be an o-decomposable Riordan graph of even order n . If G_n has at least two universal vertices, then $\det(G_n) = 0$.*

Corollary 8.5. *Let $G_n = G_n(g, f)$ be an e -decomposable Riordan graph of even order n . Then, $\det(B) = 0$ if and only if $\det(G_n) = 0$, that is, G_n is a singular graph if and only if H_n is such a graph. In particular, if $N_{H_n}(i) = N_{H_n}(j)$ for $i, j \in V_e$, $i \neq j$, then $\det(G_n) = 0$.*

Corollary 8.6. *Let $G_n = G_n(g, f)$ be an e -decomposable Riordan graph of even order n . If G_n has at least two universal vertices, then $\det(G_n) = 0$.*

The following gives an example for Corollary 8.5.

Example 8.7. *Let $G_8 = G_8\left(\frac{1}{1-z^2}, \frac{z}{1-z}\right)$. Since g is an even function, G_8 is an e -decomposable Riordan graph. The matrix B for this graph is*

$$B = \begin{bmatrix} 1 & 1 & 1 & 1 \\ 1 & 1 & 0 & 1 \\ 0 & 1 & 1 & 1 \\ 1 & 1 & 1 & 1 \end{bmatrix}.$$

Since $\mathbf{b}_1 = \mathbf{b}_4$, $\det(B) = 0$, and then, by Corollary 8.5, $\det(G_8) = 0$.

We end this section by considering determinants of Catalan graphs CG_n . Let CM_n be the adjacency matrix of CG_n . Since CG_n is a Bell type io-decomposable Riordan graph, from Theorem 12 in [3], there exists a permutation matrix P such that

$$A_n = P(CM_n)P^T = \begin{bmatrix} CM_{\lceil \frac{n}{2} \rceil} & B \\ B^T & O \end{bmatrix}. \quad (26)$$

By the binomial formula for the Catalan matrix $\left(\frac{1-\sqrt{1-4z}}{2z}, \frac{1-\sqrt{1-4z}}{2}\right)$, we have that

$$CM_n(i, j) \equiv C(i-2, j-1) = \frac{j}{i-1} \binom{2i-j-3}{i-j-1}, \quad i > j \geq 1. \quad (27)$$

Let \mathbf{b}_j be the j th row of the submatrix B .

Lemma 8.8. *For the block B in the matrix A_n , we have*

$$\mathbf{b}_2 \equiv \mathbf{b}_3. \quad (28)$$

Proof: To prove (28), it suffices to prove that

$$CM_n(3, j) = CM_n(5, j) \quad (29)$$

for each even j , $2 \leq j \leq n$. For $j \in \{2, 4\}$, we have $CM_n(3, j) = CM_n(5, j) = 1$. Suppose that $j = 2k$ where $3 \leq k \leq 2 \lfloor \frac{n}{2} \rfloor$. By (27), we have

$$C(2k-2, 2) = \frac{3(2k-3)(4k-7)}{5(k-2)(2k-5)} C(2k-2, 4) \equiv \frac{C(2k-2, 4)}{k-2}. \quad (30)$$

If k is odd, then by (27) and (30),

$$CM_n(2k, 3) \equiv C(2k-2, 2) \equiv \frac{C(2k-2, 4)}{k-2} \equiv CM_n(2k, 5),$$

which gives us (29). Otherwise k is even. With $C_n = \frac{1}{n+1} \binom{2n}{n}$ denoting the n th Catalan number,

we have

$$\begin{aligned}
C(2k-2, 4) &= \frac{5}{2k-1} \binom{4k-8}{2k-6} \\
&= \frac{5}{s+3} \binom{2s}{s-2} \quad (\text{letting } s := 2k-4) \\
&= \frac{5(s-1)s}{(s+1)(s+2)(s+3)} \binom{2s}{s} \\
&= \frac{5(s-1)s}{(s+2)(s+3)} C_s, \\
&\equiv \frac{s}{s+2} C_s = \frac{2k-4}{2k-2} C_{2k-4} \equiv (k-2) C_{2k-4} \equiv 0.
\end{aligned} \tag{31}$$

Now, by (30), (31) and Lemma 2.23, we obtain

$$C(2k-2, 2) \equiv \frac{C(2k-2, 4)}{k-2} \equiv C_{2k-4} \equiv 0$$

completing our proof. \square

Theorem 8.9. $\det(CG_{2n}) = 0$ for any $n \geq 3$.

Proof: Since the order of the matrix CM_{2n} is $2n$, the block B in (26) is a square matrix of order n . On the other hand, by Lemma 8.8 we have $\mathbf{b}_2 \equiv \mathbf{b}_3$. Thus, $\text{rank}(B) \leq n-1$ and $\det(B) = 0$. Now, we obtain the desired result from the fact that $\det(CM_{2n}) = (-1)^{2n}$. \square

Remark 8.10. Let $\Gamma = \{11, 13, 15, 23, 33, 51, 61, 63\}$. By Sage [16], we observe that for $n \in \Gamma$, $\det(CG_n) = 0$.

We conclude our paper with the following conjecture.

Conjecture 8.11. We have that $\det CG_n = 0$ if and only if $n \geq 6$ is even, or $n \in \Gamma$.

References

- [1] D. S. Bernstein, Matrix Mathematics, Princeton University Press, New York, 2005.
- [2] A. E. Brouwer, W. H. Haemers, *Spectra of graphs*, Springer, Amsterdam, 2011.
- [3] G.-S. Cheon, J.-H. Jung, S. Kitaev, S. A. Mojallal, Riordan graphs I: Structural properties, arXiv:1710.04604.
- [4] G.-S. Cheon, J. S. Kim, S. A. Mojallal, Spectral properties of Pascal graphs, *Linear and Multilinear Algebra* (2017) <https://doi.org/10.1080/03081087.2017.1356261>
- [5] D. Cvetković, P. Rowlinson, S. Simić, *An introduction to the theory of graph spectra*, Cambridge University Press, Cambridge, 2012.
- [6] K. C. Das, Sharp lower bounds on the Laplacian eigenvalues of trees, *Linear Algebra and its Applications* 384 (2004) 155–169.
- [7] K. C. Das, The Laplacian Spectrum of a Graph, *Computers and Mathematics with Applications* 48 (2004) 715–724.

- [8] K. C. Das, P. Kumar, Some new bounds on the spectral radius of graphs, *Discrete Mathematics*. 281 (2004) 149–161.
- [9] N. Deo, M. J. Quinn, Pascal Graphs and Their Properties, *The Fibonacci Quarterly* 21 (1983) 203–214.
- [10] E. Deutsch, B. E. Sagan, Congruences for Catalan and Motzkin numbers and related sequences, *J. Number Theory* 117 (2006) 191–215.
- [11] M. Fiedler, Algebraic connectivity of graphs, *Czech. Math. J.* 37 (1987) 660–670.
- [12] S. H. Ghorban, *Toeplitz graph decomposition*, *Transactions on Combinatorics* 1(4) (2012), 35–41.
- [13] D. Merlini, D.G. Rogers, R. Sprugnoli, M.C. Verri, On some alternative characterizations of Riordan arrays, *Canadian J. Mathematics* 49 (1997) 301–320.
- [14] R. Merris, Laplacian matrices of graphs: A survey, *Linear Algebra Appl.* 197, 198 (1994) 143–176 .
- [15] B. Mohar, The Laplacian Spectrum of graphs, *Graph Theory, Combinatorics and Applications* (Alavi, Y. et al., Eds.), John Wiley, New York,(1991) 871–898.
- [16] W.A. Stein, et al., Sage Mathematics Software (Version 6.8), The Sage Development Team, <http://www.sagemath.org>, 2015.
- [17] L. V. Shapiro, S. Getu, W. J. Woan, L. Woodson, The Riordan group, *Discrete Appl. Math.* 34 (1991) 229–239.
- [18] B. P. Sinha, S. Ghose, B. B. Bhattacharya, A further note on Pascal graphs, *The Fibonacci Quarterly* 24 (1986) 251–257.
- [19] H. Täubig, *Matrix Inequalities for Iterative Systems*, CRC Press, Taylor & Francis Group, Boca Raton, 2017.